

## The reactions of cyclotriphosphazene with 2-(2-hydroxyethylamino)ethanol. Spectroscopic studies of the derived products

Seung Seo Lee, Pushpak Mizar, Ozge Kucuk & Sedat Ture

To cite this article: Seung Seo Lee, Pushpak Mizar, Ozge Kucuk & Sedat Ture (2020) The reactions of cyclotriphosphazene with 2-(2-hydroxyethylamino)ethanol. Spectroscopic studies of the derived products, Phosphorus, Sulfur, and Silicon and the Related Elements, 195:6, 454-463, DOI: [10.1080/10426507.2020.1712396](https://doi.org/10.1080/10426507.2020.1712396)

To link to this article: <https://doi.org/10.1080/10426507.2020.1712396>



Published online: 11 Jan 2020.



Submit your article to this journal [↗](#)



Article views: 292



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)



# The reactions of cyclotriphosphazene with 2-(2-hydroxyethylamino)ethanol. Spectroscopic studies of the derived products

Seung Seo Lee<sup>a</sup>, Pushpak Mizar<sup>a</sup>, Ozge Kucuk<sup>b</sup>, and Sedat Ture<sup>b</sup> 

<sup>a</sup>Chemistry, Department of Natural and Environmental Sciences, Southampton University, Southampton, UK; <sup>b</sup>Department of Chemistry, Molecular Biology and Genetics, Faculty of Arts & Sciences, University of Bilecik Seyh Edebali, Bilecik, Turkey

## ABSTRACT

The reactions of cyclotriphosphazene (**1**) with 2-(2-hydroxyethylamino)-ethanol (**2**) were investigated. 2-(2-hydroxyethylamino)-ethanol (**2**) is a tri-functional reagent consisting of both aliphatic hydroxyl and the secondary amino groups and its nucleophilic substitution reactions with cyclotriphosphazene can lead to different product types; open chain, spiro, ansa, bridged and their mixtures. The reactions with one, two and three equimolar ratios of 2-(2-hydroxyethylamino)-ethanol, in the presence of NaH at 0–10 °C and at room temperature gave the following cyclotriphosphazene derivatives: one mono-spiro, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O] (**3**, 1:1, r.t.); its isomer mono-ansa (**5**, 1:1, r.t.); one dispiro, N<sub>3</sub>P<sub>3</sub>Cl<sub>2</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O]<sub>2</sub> (**4**, 1:1, r.t.); its isomer spiro-ansa (**6**, 1:2, r.t.); and one single-bridged compound with spiro substituted units, N<sub>3</sub>P<sub>3</sub>Cl<sub>3</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O]<sub>3</sub>N<sub>3</sub>P<sub>3</sub>Cl<sub>3</sub> (**7**, 1:3, at 0–10 °C); as well as single-, N<sub>3</sub>P<sub>3</sub>Cl<sub>5</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O]<sub>3</sub>N<sub>3</sub>P<sub>3</sub>Cl<sub>5</sub> (**8**, 1:2, r.t.), double-, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O]<sub>2</sub>N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub> (**9**, 1:2, r.t.), and tri-bridged, N<sub>3</sub>P<sub>3</sub>Cl<sub>3</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>-O]<sub>3</sub>N<sub>3</sub>P<sub>3</sub>Cl<sub>3</sub> (**10**, 1:3, at 0–10 °C) derivatives. Triple-bridged derivative is the major product in this system. The structures of the novel-derived compounds were characterized by TLC-MS, FT-IR, elemental analysis, <sup>1</sup>H, and <sup>31</sup>P NMR spectral.


## ARTICLE HISTORY

Received 4 November 2019  
Accepted 3 January 2020

## KEYWORDS

Cyclotriphosphazene; 2-(2-hydroxyethylamino)ethanol; spiro compounds; ansa compound; bridged derivatives

## GRAPHICAL ABSTRACT



## 1. Introduction

Due to having a variety of applications in science and technology, cyclophosphazenes are an important class of inorganic cyclic systems<sup>[1,2]</sup> and therefore, reactions of cyclotriphosphazene (**1**) with mono-, di- and polyfunctional nucleophilic reagents have been extensively studied by different research groups. The most important feature of the cyclophosphazene chemistry is that the halogen atoms attached to the phosphorus atoms in the cyclic system can be replaced by nucleophilic substitution reactions with various organic, inorganic or organometallic reagents and the compounds exhibit different physical, chemical and biological properties relative to the substituted ligands.<sup>[1–46]</sup> Cyclic phosphazenes-based derivatives have wide range of properties and applications such as electrical conductivity,<sup>[24]</sup> liquid crystals,<sup>[42,44–47]</sup> chemosensors<sup>[28]</sup> and biologically active materials.<sup>[14–20]</sup> In recent years, many studies have been performed on cyclophosphazene derivatives to determine cytotoxic and antimicrobial activities and their interactions with DNA.<sup>[20–23,26,27,29–35]</sup>

The reactions of cyclotriphosphazene (**1**) with tri-functional nucleophilic reagents have been previously studied.<sup>[48–52]</sup> The reactions with spermidine (as a triamine) gave only one product, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>NH[(CH<sub>2</sub>)<sub>3</sub>NH(CH<sub>2</sub>)<sub>4</sub>NH]<sub>2</sub>N<sub>3</sub>P<sub>3</sub>Cl<sub>5</sub>. This was confirmed by single crystal X-ray diffraction to have a 6-membered spiro ring, involving one N<sub>3</sub>P<sub>3</sub> moiety and a bridge to a second such unit.<sup>[48,49]</sup>

On the other hand, from the reactions of cyclotriphosphazene with glycerol (as a triol) two isomeric products were reported to be in the form of an oil and a crystalline material.<sup>[50–52]</sup> While the oily fraction (as the major product) was assigned the 5-membered spiro structure, the crystalline one was identified as the 6-membered spiro structure, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>[O<sub>3</sub>C<sub>3</sub>H<sub>6</sub>].<sup>[50–52]</sup>

We therefore wished to expand and clarify the reactivity and the substitution patterns of cyclotriphosphazene (**1**) with tri-functional nucleophilic reagents. And, in this study, hexachlorocyclotriphosphazatriene (**1**) was reacted with 2-(2-hydroxyethylamino)-ethanol (**2**, DEA) and the novel




Figure 1. Structures of the synthesized cyclotriphosphazene derivatives (3–10).

secondary amine containing spiro-, ansa- and bridged-cyclotriphosphazene derivatives (3–10, Figure 1) were synthesized. The structural characterization of the compounds was performed by using elemental analyses, MS,  $^1H$ ,  $^{31}P$ - $\{^1H\}$ , and  $^1H\{^{31}P\}$  NMR spectral data.

## 2. Results and discussion

The reactions of cyclotriphosphazene,  $N_3P_3Cl_6$  (1), with 2-(2-hydroxyethylamino)ethanol,  $HO-CH_2-CH_2-NH-CH_2-CH_2-OH$  (2), in excess of NaH, at 0–10 °C and at room temperature, and in 1:1, 1:2 and 1:3 molar ratios gave the following novel *geminal*- and *non-geminal*-substituted cyclophosphazene derivatives: one mono *spiro*,  $N_3P_3Cl_4[O-(CH_2)_2-NH-(CH_2)_2O]$  (3, 1:1, r.t. 27%); its ansa isomer (5, 1:1, r.t. 33%); one dispiro,  $N_3P_3Cl_2[O-(CH_2)_2-NH-(CH_2)_2O]_2$  (4, 1:1, r.t. 19%); its spiro-ansa isomer (6, 1:2, r.t. 25%); and one single-bridged compound with spiro substituted units,  $N_3P_3Cl_3[O-(CH_2)_2-NH-(CH_2)_2-O]_3$  (7, 1:3, at 0–10 °C, 14%); as well as single-,  $N_3P_3Cl_5[O-(CH_2)_2-NH-(CH_2)_2O]_2$  (8, 1:2, r.t. 23%); double-,  $N_3P_3Cl_4[O-(CH_2)_2-NH-(CH_2)_2O]_2$  (9, 1:2, r.t. 32%); and triple-bridged  $N_3P_3Cl_3[O-(CH_2)_2-NH-(CH_2)_2O]_3$  (10, 1:3, at 0–10 °C, 45%) products. The structures of the compounds were verified by elemental analysis, TLC-MS,  $^1H$  and  $^{31}P$  NMR spectroscopy and the


results are provided as part of the analytical data in the section on synthesis. The structures of the derived compounds (3–10) are exhibited in Figure 1.

### 2.1. $^{31}P$ NMR spectra

At first, the proton-decoupled  $^{31}P$  NMR spectra of the reaction mixtures were measured to determine which product types were formed and their relative proportions. Then, proton-decoupled and proton-coupled  $^{31}P$  NMR spectra were used to characterize the structures of the derived compounds. The new synthesized cyclophosphazene derivatives (3–10) contain  $\equiv PCl_2$ ,  $\equiv P(OR)_2$  and  $\equiv P(OR)Cl$  groups and the  $^{31}P$  NMR spectra are observed as  $A_2X$  (or  $A_2B$ ) and  $AMX$  spin types,<sup>[2,3,11,12,53]</sup> which can be readily assigned by consideration of signal intensities, chemical shifts and coupling patterns. Selected  $^{31}P$  NMR chemical shifts and coupling constants of compounds (3–10) are summarized in Table 1. Within the  $A_2X$  (or  $AB_2$ ) spin systems, the proton-coupled  $^{31}P$  NMR spectra suggest that the  $A_2$  parts of the spectra arise from  $\equiv PCl_2$  (in compounds 3, 8);  $\equiv P(OR)Cl$  (in compounds 5, 6 and 9) and  $\equiv P(OR)OR$  (in compound 4) groups, giving rise to doublet structures and the X parts of the spectra arise from  $\equiv P(OR)_2$  (in compounds 3 and 6),  $\equiv PCl_2$  (in compounds 4, 5 and 9) and the  $\equiv P(OR)Cl$  (in compound 8) groups, giving rise to triplet structures.

**Table 1.** Selected  $^{31}\text{P}$  NMR parameters for compounds 3–10<sup>a</sup>.

Compound	$\delta\text{P}(\text{OR})_2^b$	$\delta\text{P}(\text{OR})\text{Cl}^b$	$^2J_{\text{P}(\text{OR})_2-\text{P}(\text{OR})\text{Cl}}^c$	$^2J_{\text{P}(\text{OR})_2-\text{P}(\text{OR})\text{Cl}}^c$
(3)	22.40	14.60	68.70	
(4)	28.10	13.20		62.70
(5)	25.80	16.50		76.30
(6)		15.40		83.40 <sup>d</sup>
(7)	29.40	48.80	1.60	<sup>e</sup>
(8) single-bridged	26.80	15.50		65.75
(8) double-bridged	27.50	20.30		64.30
(10) triple-bridged		8.62		


<sup>a</sup>In  $\text{CDCl}_3$  (with respect to 85% phosphoric acid external reference) at 162 and 202.38 MHz.<sup>b</sup>In ppm.<sup>c</sup>In Hz.<sup>d</sup> $^2J_{\text{P}(\text{OR})_2-\text{P}(\text{OR})\text{Cl}}$ .<sup>e</sup>Unresolved spectrum.**Figure 2.** (a)  $^{31}\text{P}\{-^1\text{H}\}$  and (b)  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of compound 3, in  $\text{CDCl}_3$  at 162.00 MHz, (room temperature), referenced to external 85%  $\text{H}_3\text{PO}_4$ .

The proton-coupled  $^{31}\text{P}$  NMR spectra allow identification of the lines due to the  $\equiv\text{P}(\text{OR})_2$  and  $\equiv\text{P}(\text{OR})\text{Cl}$  groups, where each group splits into further lines. Proton coupling experiments as well as comparison with the earlier reported studies with difunctional and tri-functional reagents,<sup>[5–7,10,48–52]</sup> allow unambiguous assignments of the structures. The proton-coupled and decoupled  $^{31}\text{P}$  NMR spectra of compounds 3 and 4 are exhibited in Figure 2 and 3 respectively.

In the triple-bridged cyclophosphazene derivative (10), the phosphorus atoms were substituted with  $\text{P}(\text{OR})\text{Cl}$

groups and therefore, all the phosphorus atoms are chemically and magnetically in the same environment and the  $\equiv\text{P}(\text{OR})\text{Cl}$  groups are symmetrically located in the molecule. Therefore, the  $^{31}\text{P}$  NMR spectrum gave rise to a single line ( $A_6$  spin system), whereas all the other cyclophosphazene derivatives, as explained above, contain  $\equiv\text{PCl}_2$ ,  $\equiv\text{P}(\text{OR})_2$  or  $\equiv\text{P}(\text{OR})\text{Cl}$  groups and the  $^{31}\text{P}$  NMR spectra were observed as  $A_2X$  (or  $A_2B$ ) spin types.

The  $^{31}\text{P}$  NMR spectrum of the ansa compound (5) is of the  $A_2X$ , type with a doublet for the  $\equiv\text{P}(\text{OR})\text{Cl}$  and a triplet



**Figure 3.** (a)  $^{31}\text{P}\{-^1\text{H}\}$  and (b)  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of compound 4, in  $\text{CDCl}_3$  at 162.00 MHz, (room temperature), referenced to external 85%  $\text{H}_3\text{PO}_4$ .

for the  $\equiv\text{PCl}_2$  groups at 16.80 and 25.80 ppm respectively.  $^{31}\text{P}$  NMR proton coupling affects the  $\text{A}_2$  ( $\equiv\text{P(OR)Cl}$ ) part of the spectrum by showing further splitting on it. Cylophosphazene derivative 7 is a compound formed by interconnecting two mono-spiro structures with a single-bridged link and has an AMX spin system comprising both the spiro [ $\equiv\text{P(OR)}_2$ ], the bridged [ $\equiv\text{P(OR)Cl}$ ] and the  $\equiv\text{PCl}_2$  groups. Therefore, the phosphorus atoms situated in three different magnetic environments. Both phosphorus atoms within each group; the spiro [ $\equiv\text{P(OR)}_2$ , at 48 ppm] parts, the  $\equiv\text{PCl}_2$  parts (at 29.40 ppm) and the  $\equiv\text{P(OR)Cl}$  moieties (at 1.60 ppm) are identical to each other and symmetrically located in the molecule. But those phosphorus atoms in three different groups are not in the same magnetic environment and therefore, it is of the AMX spin type.

$^{31}\text{P}$  NMR proton-coupled spectrum gave rise to further splitting at 48.80 ppm and at 1.60 ppm for the spiro,  $\equiv\text{P(OR)}_2$  and the bridged,  $\equiv\text{P(OR)Cl}$  moieties. Phosphorus proton decoupled and coupled NMR spectra of compound 7 are shown in Figures 4 and 5 respectively.

## 2.2. $^1\text{H}$ NMR data

In this system, the methylene and the amino-group protons occur in three different chemical environments depending on whether they are situated,  $\alpha$ - and  $\beta$ - with respect to the oxygen atoms and  $\text{NH}$  protons. The  $^1\text{H}$  NMR spectra of compounds (4–7) gave very complex multiplets with some overlapping signals and similar chemical shifts for the




Figure 4.  $^{31}\text{P}\{-^1\text{H}\}$  NMR spectrum of compound 7, in  $\text{CDCl}_3$  at 162.00 MHz, (room temperature), referenced to external 85%  $\text{H}_3\text{PO}_4$ .




Figure 5.  $^{31}\text{P}\{^1\text{H}\}$  coupled NMR spectrum of compound 7, in  $\text{CDCl}_3$  at 162.00 MHz, (room temperature), referenced to external 85%  $\text{H}_3\text{PO}_4$ .

$\text{POCH}_2$  and the  $\text{POCCH}_2$  protons. We observed much more complex splitting patterns for compounds (6) and (7) for the  $\text{POCH}_2$  and  $\text{POCCH}_2$  methylene protons, suggesting that the  $\alpha$ - and  $\beta$ -methylene protons are not sufficiently resolved.

The  $\alpha$ - and the  $\beta$ -methylene protons of the mono-spiro (3) compound within each methylene group are in the same magnetic environment and therefore, the mono-spiro derivative, has a relatively simple proton spectrum, giving a six line spectrum for the methylene protons ( $\text{POCH}_2$ ) at 4.46 ppm, from coupling with the neighboring  $\beta$ -methylene protons and the phosphorus nucleus. On the other hand the  $\beta$ -methylene protons ( $\text{POCH}_2\text{-CH}_2\text{-NH-}$ ) give two quintets spectra at 1.62 ppm due to coupling with the two  $\alpha$ -, N-H protons and with the phosphorus nucleus. The  $^1\text{H}$  NMR spectra of compounds (3) and (8) are presented in Figures 6 and 7. The  $\alpha$ - and the  $\beta$ -methylene protons are not chemically and magnetically equivalent in compounds 4 and 5, in

the ansa and in the spiro parts of compound 6, and in the spiro and in the bridging units of compound 7. For this reason, at 400 MHz, very complex and unresolved spectra were observed for the  $\alpha$ - and the  $\beta$ -methylene protons. Therefore, detailed consideration for the  $^1\text{H}$  NMR spectra of the compounds (4–7) are not possible. However, we tried to simplify these spectra with comparing to our earlier reported studies based on difunctional and tri-functional reagents.<sup>[3–7,10,48–52]</sup> Selected  $^1\text{H}$  NMR chemical shifts and coupling constants for the  $\alpha$ - and the  $\beta$ -methylene protons are summarized in Table 2. In the ansa moiety (5) the two  $\alpha$ - and the two  $\beta$ -methylene groups are equivalent, but the two methylene protons within each methylene group are nonequivalent as one faces toward, the other away from, the  $\text{N}_3\text{P}_3$  ring and its spiro group. In the spiro part of the spiro-ansa compound (6) the  $\alpha$ -methylene protons as well in the  $\beta$ -methylene protons are nonequivalent as the group above and




Figure 6.  $^1\text{H}$  NMR spectrum of compound (3), at room temperature, in  $\text{CDCl}_3$  (TMS as internal reference) and at 399.95 MHz.




Figure 7.  $^1\text{H}$  NMR spectrum of compound (8), at room temperature, in  $\text{CDCl}_3$  (TMS as internal reference) and at 399.95 MHz.

below the phosphazene ring are in different chemical environment, those above the ring seeing the ansa group, those below the chlorine atoms. However, within each methylene group the two protons are equivalent. By contrast, in the ansa moiety the two  $\alpha$ -methylene groups are equivalent. The same applies to the  $\beta$ -methylene groups, but all the methylene groups,  $\alpha$ - and  $\beta$ -methylene protons have nonequivalent protons. The  $^1\text{H}$  NMR spectrum of the bis-spiro derivative (4) is by far the most complex and also the most interesting, where two  $\text{POCH}_2$  and two  $\text{CCH}_2$  methylene groups are

equivalent, but within each methylene group the two protons are nonequivalent due to their being part of a cyclic moiety and therefore the two protons of each methylene group see a different environment. In dispiro structures, the two spiro rings are equivalent as are the  $\alpha$ - and the  $\beta$ -methylene groups above and below the rings. However, because each spiro group is flanked on one side by two chlorine atoms and on the other side by another spiro group, the two protons within each methylene group are nonequivalent. This nonequivalence was quite noticeable for

the  $\alpha$ -methylene protons and even more pronounced for the  $\beta$ -methylene protons. If such a nonequivalence is observed all the methylene groups could in principle give rise to AB quartets with further coupling to neighboring nuclei. Long-range virtual coupling with the two equivalent phosphorus nuclei will make a triplet of quartets for the  $\alpha$ -methylene protons and coupling with the  $\beta$ -methylene protons might, depending on Karplus relationship,<sup>[54,55]</sup> give rise to 36 or 48 lines.

### 3. Experimental

#### 3.1. Materials

Reagent grade solvents were used throughout the work, benzene, light petroleum (b.p. 40–60°C), petroleum ether, anhydrous diethyl ether, acetonitrile, butanol, n-hexane (>96%), dichloromethane (>99.0%), chloroform, acetone, THF, acetone (Sigma Aldrich). THF was distilled over a sodium-potassium alloy under an argon atmosphere. CDCl<sub>3</sub>, deuteriated solvent for NMR spectroscopy (Sigma Aldrich), Silica gel (60, 0.063–0.200 mm Merck) was used for column chromatography, Kieselgel 60° 254 (silica gel) precoated TLC plates (Merck). The following materials were also obtained from Sigma Aldrich Chemicals: Phosphonitrilic trimer (purified by fractional crystallization from hexane), ninhydrine (0.5% w/v), 2-(2-hydroxyethylamino)ethanol (from Southampton University Laboratory), NaH (60% dispersion in mineral oil, which was removed by washing with dry n-heptane followed by decantation).

#### 3.2. Methods

All reactions were monitored using Kieselgel 600 254 (silica gel) precoated TLC plates and sprayed with ninhydrine (0.5% w/v) in butanol solution, and developed at approximately

130 °C. UV for TLC measurements. Required separations of mixtures were carried out by crystallization techniques, TLC and by column chromatography using Kieselgel 60. (Merck 60, 0.063–0.200 mm; for 2 g crude mixture, 100 g silica gel was used in a column of 2.5 cm in diameter and 90 cm in length). Melting points were determined on a Hot Stage Microscopy and hot stage connected to a FP 800 central processor both fitted with a polarizing microscope at Southampton University. FT-IR spectra were recorded in Perkin Elmer BX II FT model spectrometer with a number of scans at 4 cm<sup>-1</sup> resolution in the range 4000–350 cm<sup>-1</sup>. Elemental analyses were obtained using a ThermoFinnigan Flash 1112 Instrument. <sup>1</sup>H NMR spectra were recorded using a Bruker AVIIIHD 400 MHz spectrometer (operating at 399.5 MHz, at Southampton University). Samples were dissolved in CDCl<sub>3</sub> and placed in 5 mm NMR tubes. Measurements were carried out using a CDCl<sub>3</sub> lock, TMS as internal reference, and sample concentrations of 15–20 mg cm<sup>3</sup>. <sup>13</sup>C NMR spectra were recorded using a Bruker AVIIIHD 400 MHz spectrometer (operating at 100.63 MHz, at Southampton University), TMS was used as an internal

reference. <sup>31</sup>P NMR spectra were recorded using a Bruker AVIIIHD 400 MHz spectrometer (operating at 161.97 MHz, at Southampton University); Measurements were carried out using a CDCl<sub>3</sub> lock and H<sub>3</sub>PO<sub>4</sub> (85%) was used as an external reference. Mass spectra were recorded using a TLC/MS (obtained by a Bruker MicroTOF LC/MS spectrometer using electro spray ionization (ESI) method). The NMR data may be found in Tables 1 and 2.

#### 3.3. The reactions of hexachlorocyclotriphosphazatriene (1) with 2-(2-hydroxyethylamino)ethanol (2)

(a) One equivalent of compound 2: Using an excess of NaH, in a mixture of THF/ethanol (3:1) at room temperature. Synthesis of 2,2-mono-spiro-2,2'-iminodiethoxy-4,4,6,6-tetrachloro-cyclotriphosphazatriene (3); 2,2,4,4-bis-spiro-2,2'-iminodiethoxy-6,6-dichloro-cyclotriphosphazatriene (4); and 2,4-ansa-2,2'-iminodiethoxy-2,4,6,6-tetrachloro-cyclotriphosphazatriene (5).

Cyclotriphosphazene (1, 2 g, 0.006 mol) and 2-(2-hydroxyethylamino)ethanol (2, 0.63 g, 0.006 mmol) were dissolved in dry THF (100 mL) in a 250 mL three-necked round-bottom flask. This mixture was stirred for 30 min at room temperature then two equivalents of NaH (0.29 g, 0.012 mol) in THF (30 mL) was added over a period of 30–40 min to this stirred solution under an argon atmosphere. The mixture was stirred (36 h) at room temperature until TLC indicated the completion of the reaction. The reaction mixture was filtered to remove sodium chloride and any other insoluble materials. Then the reaction mixture was followed on TLC silica gel plates using dichloromethane-hexane (3:1) as the eluent. The solvent was removed under reduced pressure and the resulting colorless solid was subjected to column chromatography using the mentioned solvent system as the mobile phase. Products were recrystallized from benzene: hexane (1:3) containing a few drops of light petroleum (b.p. 40–60 °C). Three main fractions were synthesized as (i) the mono spiro derivative, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>O] (3), recrystallized from dichloromethane-hexane (1:1.5) by adding a few drops of light petroleum (b.p. 40–60 °C), yield 0.5 g, 27%, m.p. 128–129 °C. Anal. Calc. for N<sub>4</sub>P<sub>3</sub>Cl<sub>4</sub>C<sub>4</sub>H<sub>9</sub>O<sub>2</sub>: C, 12.64; H, 2.38; N, 14.74, M, 379.94. Found: C, 12.63; H, 2.36; N, 14.52, M<sup>+</sup>, 380.69. <sup>1</sup>H NMR (CDCl<sub>3</sub>),  $\delta$ POCH<sub>2</sub>: 4.46,  $\delta$ POCCH<sub>2</sub>: 1.62,  $\delta$ PNH: 7.25, <sup>3</sup>J<sub>(PH)</sub>: 14.40 Hz. <sup>31</sup>P NMR (CDCl<sub>3</sub>):  $\delta$ : 22.40 (PCL<sub>2</sub>),  $\delta$  = 14.60 [P(OR)<sub>2</sub>], <sup>2</sup>J<sub>(PP)</sub>: 68.70 Hz. (ii) The second product was identified as the dispiro derivative, N<sub>3</sub>P<sub>3</sub>Cl<sub>2</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>O]<sub>2</sub> (4), 0.38 g, 19%, m.p. 174–176 °C. Anal. Calc. for N<sub>5</sub>P<sub>3</sub>Cl<sub>2</sub>C<sub>8</sub>H<sub>18</sub>O<sub>4</sub>: C, 23.32; H, 4.40; N, 16.99, M, 412.17. Found: C, 22.99; H, 4.43; N, 16.95, M<sup>+</sup>, 413.09. <sup>1</sup>H NMR (CDCl<sub>3</sub>);  $\delta$ POCH<sub>2</sub>: 4.46/4.50,  $\delta$ POCCH<sub>2</sub>: 1.66,  $\delta$ PNH: 7.36, <sup>3</sup>J<sub>(PH)</sub>: 18.20. <sup>31</sup>P NMR (CDCl<sub>3</sub>);  $\delta$ : 28.10 (PCL<sub>2</sub>),  $\delta$ : 13.20 [P(OR)<sub>2</sub>], <sup>2</sup>J<sub>(PP)</sub>: 62.70 Hz. (iii) The third product was identified as the mono-ansa derivative, N<sub>3</sub>P<sub>3</sub>Cl<sub>4</sub>[O-(CH<sub>2</sub>)<sub>2</sub>-NH-(CH<sub>2</sub>)<sub>2</sub>O] (5), 0.57 g, 33%, m.p., an oil. Anal. Calc. for N<sub>4</sub>P<sub>3</sub>Cl<sub>4</sub>C<sub>4</sub>H<sub>9</sub>O<sub>2</sub>: C, 12.64; H, 2.38; N, 14.74, M, 379.94. Found: C, 12.60; H, 2.40; N, 14.52, M<sup>+</sup>, 380.69. <sup>1</sup>H NMR (CDCl<sub>3</sub>),  $\delta$ POCH<sub>2</sub>: 4.44–4.58,  $\delta$ POCCH<sub>2</sub>:

**Table 2.** Selected  $^1\text{H}$  NMR parameters for compounds **3–10**<sup>a</sup>.

Compound	$\delta\text{POCH}_2^b$	$\delta\text{PCCH}_2^b$	$\delta\text{NH}^b$	$^3\text{J}(\text{PH})^c$
(3) mono-spiro	4.46	1.62	7.25	14.40
(4) bis-spiro	4.50/4.46	1.66	7.36	18.20
(5) mono-ansa	4.58/4.44	1.50	7.50	21.40
(6) spiro-ansa				
spiro part	4.49	1.66	7.26	<sup>d</sup>
ansa part	4.56/4.61	1.52	7.52	<sup>d</sup>
(7) spiro-bino	4.50/4.43		6.98/7.15/7.56	<sup>d</sup>
(8) single-bridged	4.32	1.45	7.56	12.60
(9) double-bridged	4.56	1.48	7.56	12.80
(10) triple-bridged	4.42	1.64	7.70	13.43

<sup>a</sup>In  $\text{CDCl}_3$  (referenced to internal TMS internal) at 399.95 MHz. (room temperature).<sup>b</sup>In ppm.<sup>c</sup>Hz.<sup>d</sup>Unresolved spectrum.

1.50,  $\delta\text{PNH}$ : 7.50,  $^3\text{J}(\text{PH})$ : 21.40 Hz.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$ : 25.80 ( $\text{PCl}_2$ ),  $\delta$ : 16.50 [ $\text{P}(\text{OR})_2$ ],  $^2\text{J}(\text{PP})$ : 76.30 Hz.

(b) Synthesis of 2,4-ansa-6,6-spiro-2,2'-iminodiethoxy-2,4-dichloro-cyclotriphosphazatriene (**6**); 2,2,4,4,6,6-trispiro-2,2'-iminodiethoxy-cyclotriphosphazatriene (**7**); and 2,2,4,4-bino-2,2'-iminodiethoxy-2,2',4,4',6,6',6'-octachloro-cyclotriphosphazatriene (**8**): Two equivalents of

2-(2-hydroxyethylamino) ethanol (**2**, 1.26 g, 0.012 mol) and cyclotriphosphazene (**1**, 2 g, 0.006 mol) were dissolved in a mixture of THF/ethanol (3:1, 120 mL) in a 250 mL three-necked round-bottomed flask under an argon atmosphere and the reaction mixture was cooled to 0 °C in ice-bath. Then four equivalents of NaH (0.58 g, 0.024 mol) were dissolved in THF (15 mL) and added dropwise into the stirred solution under an argon atmosphere. The reaction mixture was then stirred at room temperature for about 48 h. The reaction was followed on TLC using dichloromethane-hexane (3:1) as the eluent for completion of the reaction. The reaction mixture was filtered to remove the NaCl salts and some other insoluble sticky materials. Together with the starting material (**1**), four spots were observed on TLC, using dichloromethane-hexane (2:1) as the mobile phase. The solvents were removed from the reaction mixture under reduced pressure and the resulting white-colored solids were subjected to column chromatography using the same solvent system as described above. Products were recrystallized from benzene:hexane (1:3) containing a few drops of light petroleum (b.p. 40–60 °C). (i) The first product was identified as the single-bridged derivative,  $\text{N}_3\text{P}_3\text{Cl}_5[\text{O}-(\text{CH}_2)_2-\text{NH}-(\text{CH}_2)_2\text{O}]\text{N}_3\text{P}_3\text{Cl}_5$  (**8**), yield 0.30 g, 23%, m.p. an oil. Anal. Calc. for  $\text{N}_7\text{P}_6\text{Cl}_{10}\text{C}_4\text{H}_9\text{O}_2$ : C, 6.60; H, 1.25; N, 13.47, M, 727.66. Found: C, 6.58; H, 1.29; N, 13.46,  $\text{M}^+$ , 728.59.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ );  $\delta\text{POCH}_2$ : 4.32,  $\delta\text{POCCH}_2$ : 1.45,  $\delta\text{PH}$ : 7.56,  $^3\text{J}(\text{PH})$ : 12.60 Hz.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$ : 26.80 ( $\text{PCl}_2$ ),  $\delta$ : 15.50 [ $\text{P}(\text{OR})\text{Cl}$ ],  $^2\text{J}(\text{PP})$ : 65.75 Hz. (ii) The second product was identified as the spiro-ansa derivative,  $\text{N}_3\text{P}_3\text{Cl}_2[\text{O}-(\text{CH}_2)_2-\text{NH}-(\text{CH}_2)_2\text{O}]_2$  (**6**), yield 0.32 g, 25%, m.p. 162–164 °C. Anal. Calc. for  $\text{N}_5\text{P}_3\text{Cl}_2\text{C}_8\text{H}_{18}\text{O}_4$ : C, 23.32; H, 4.40; N, 16.99, M, 412.17. Found: C, 23.21; H, 4.46; N, 16.94,  $\text{M}^+$ , 413.17.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ); spiro part;  $\delta\text{POCH}_2$ : 4.49,  $\delta\text{POCCH}_2$ : 1.66,  $\delta\text{PH}$ : 7.26; ansa part;  $\delta\text{POCH}_2$ : 4.56/4.61,  $\delta\text{POCCH}_2$ : 1.52,  $\delta\text{PH}$ : 7.52,  $^3\text{J}(\text{PH})$ : unresolved spectrum.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$ : 15.40 [ $\text{P}(\text{OR})_2$ ],  $\delta$ : 28.20 [ $\text{P}(\text{OR})\text{Cl}$ ],  $^2\text{J}(\text{PP})$ : 83.40 Hz.

(iii) The third product was identified as the double-bridged derivative,  $\text{N}_3\text{P}_3\text{Cl}_4[\text{O}-(\text{CH}_2)_2-\text{NH}-(\text{CH}_2)_2\text{O}]_2\text{N}_3\text{P}_3\text{Cl}_4$  (**9**), yield 0.52 g, 32%, m.p. 242–244 °C. Anal. Calc. for  $\text{N}_8\text{P}_6\text{Cl}_8\text{C}_8\text{H}_{18}\text{O}_4$ : C, 12.64; H, 2.39; N, 14.75; M, 759.89. Found: C, 12.63; H, 2.41; N, 14.73,  $\text{M}^+$ , 760.73.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ );  $\delta\text{POCH}_2$ : 4.56,  $\delta\text{POCCH}_2$ : 1.48,  $\delta\text{PH}$ : 7.56,  $^3\text{J}(\text{PH})$ : 12.80.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$ : 27.50 ( $\text{PCl}_2$ ),  $\delta$ : 20.30 [ $\text{P}(\text{OR})\text{Cl}$ ],  $^2\text{J}(\text{PP})$ : 64.30 Hz.

(c) Synthesis of 2,2,2',2'-spiro-2,2'-iminodiethoxy-4,4'-bino-2,2'-iminodiethoxy-4,4',6,6',6'-hexachloro-cyclotriphosphazatriene: Three equivalents of compound **2**, reaction procedure as for (**b**), at 0–10 °C. (i) The first product was identified as the triple-bridged derivative,  $\text{N}_3\text{P}_3\text{Cl}_3[\text{O}-(\text{CH}_2)_2-\text{NH}-(\text{CH}_2)_2\text{O}]_3$ ,  $\text{N}_3\text{P}_3\text{Cl}_3$  (**10**), 0.45 g, 30%, m.p. 260–263 °C. Anal. Calc. for  $\text{N}_9\text{P}_6\text{Cl}_6\text{C}_{12}\text{H}_{27}\text{O}_6$ : C, 18.19; H, 3.43; N, 15.91; M, 792.1. Found: C, 18.17; H, 3.45; N, 15.89;  $\text{M}^+$ , 793.07.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta\text{POCH}_2$ : 4.42,  $\delta\text{POCCH}_2$ : 1.64,  $\delta\text{PH}$ : 7.70,  $^3\text{J}(\text{PH})$ : 13.43.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$  = 8.62 [ $\text{P}(\text{OR})\text{Cl}$ ]. (ii) The second product was identified as the mono-spiro with a singly-bridged part,  $\text{N}_3\text{P}_3\text{Cl}_3[\text{O}-(\text{CH}_2)_2-\text{NH}-(\text{CH}_2)_2\text{O}]\text{N}_3\text{P}_3\text{Cl}_3$  (**7**), 0.18 g, 14%, m.p. an oil. Anal. Calc. for  $\text{N}_9\text{P}_6\text{Cl}_6\text{C}_{12}\text{H}_{27}\text{O}_6$ : C, 18.19; H, 3.43; N, 15.91; M, 792.10. Found: C, 18.17; H, 3.47; N, 15.92,  $\text{M}^+$ , 793.07.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ );  $\delta\text{POCH}_2$ : 4.43/4.50,  $\delta\text{POCCH}_2$ : unresolved complex spectrum,  $\delta\text{PNH}$ : 6.98/7.15/7.56,  $^3\text{J}(\text{PH})$ : unresolved spectrum.  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ );  $\delta$ : 29.40 ( $\text{PCl}_2$ ),  $\delta$ : 1.60 [ $\text{P}(\text{OR})_2$ ],  $\delta$ : 48.80 [ $\text{P}(\text{OR})\text{Cl}$ ],  $^2\text{J}(\text{PP})$ : unresolved.

## 4. Conclusion

In this study, spiro, ansa, bridged compounds as well as combination of these (**3–10**) were obtained by interacting hexachlorocyclotriphosphazatriene (**1**) with 2-(2-hydroxyethylamino)ethanol (**2**). The reactions of this tri-functional nucleophilic reagent (**2**) with compound **1** gave both the *geminal* and the *non-geminal* substitution patterns at the  $\text{PCl}_2$  moieties. The mono-spiro-bino substituted cyclophosphazene derivative (**7**) is the most interesting product for its regioselectivity and spectroscopic properties. The novel synthesized cyclotriphosphazene derivatives (**3–10**) were characterized by elemental analysis, TLC-MS,  $^1\text{H}$ , and  $^{31}\text{P}$  NMR spectral data.

## Acknowledgement

We are indebted to the School of Chemistry/Southampton University for providing a fully facilitated academic visitor's research placement and providing great opportunity to obtain spectroscopic data.

## ORCID

Sedat Ture  <http://orcid.org/0000-0001-8637-5580>

## References

- [1] Gleria, M.; DeJaeger, R. (Eds.) *Applicative Aspects of Cyclophosphazenes*; Nova Science: New York, **2004**; 1048.
- [2] (a) Allcock, H. R. *Phosphorus-Nitrogen Compounds. 1st Addition, Cyclic, Linear, and High Polymeric Systems*; Academic Press: New York and London, **1972**; pp 1–512. ISBN:

9780323147514. DOI: [10.1016/B978-0-12-050560-9.X5001-3](https://doi.org/10.1016/B978-0-12-050560-9.X5001-3). (b) Allen, C. W. Regio- and Stereochemical Control in Substitution Reactions of Cyclophosphazenes. *Chem. Rev.* **1991**, *91*, 119–135. DOI: [10.1021/cr00002a002](https://doi.org/10.1021/cr00002a002).
- [3] Shaw, R. A. The Reactions of Phosphazenes with Difunctional and Polyfunctional Nucleophilic Reagents. *Phosphorus Sulfur Silicon Rel. Elem.* **1989**, *45*, 103–136. DOI: [10.1080/10426508908046081](https://doi.org/10.1080/10426508908046081).
- [4] Guerch, G.; Labarre, J. F.; Lahana, R.; Roques, R.; Sournies, F. An Answer to the Spiro versus Ansa Dilemma in Cyclophosphazenes. *J. Mol. Struct.* **1983**, *99*, 275–282. DOI: [10.1016/0022-2860\(83\)90030-3](https://doi.org/10.1016/0022-2860(83)90030-3).
- [5] Castera, P.; Faucher, J. P.; Guerch, G.; Lahana, R.; Mahmoud, A.; Sournies, F.; Labarre, J. F. An Answer to the SPIRO versus ANSA Dilemma in Cyclophosphazenes. Part VII. Neither SPIRO nor ANSA: The BINO Dicyclopentaphosphazenes,  $N_3P_3Cl_5$ –[HN–(CH<sub>2</sub>)<sub>n</sub>–NH]Cl<sub>5</sub>P<sub>3</sub>N<sub>3</sub>. *Inorg. Chim. Acta* **1985**, *108*, 29–33. DOI: [10.1016/S0020-1693\(00\)84319-4](https://doi.org/10.1016/S0020-1693(00)84319-4).
- [6] Coles, S. J.; Davies, D. B.; Eaton, R. J.; Hursthouse, M. B.; Kiliç, A.; Mayer, T. A.; Shaw, R. A.; Yenilmez, G. Chiral Configurations of Spermine-Bridged Cyclophosphazatrienes. *J. Chem. Soc. Dalton Trans.* **2002**, *3*, 365–370. DOI: [10.1039/b104973a](https://doi.org/10.1039/b104973a).
- [7] Murr, E. N.; Lahana, R.; Labarre, J. F. The Dispiro  $N_3P_3Cl_2$  [HN–(CH<sub>2</sub>)<sub>3,4</sub>–NH]<sub>2</sub> and Trispiro  $N_3P_3$ [HN–(CH<sub>2</sub>)<sub>3</sub>–NH]<sub>3</sub> Derivatives. *J. Mol. Struct.* **1984**, *117*, 73–85. DOI: [10.1016/0022-2860\(84\)87244-0](https://doi.org/10.1016/0022-2860(84)87244-0).
- [8] Allcock, H. R.; Diefenbach, U.; Pucher, S. R. New Mono- and Tri-Spirocyclophosphazenes from the Reaction of (NPCl<sub>2</sub>)<sub>3</sub> with Aromatic Ortho Dinucleophiles. *Inorg. Chem.* **1994**, *33*, 3091–3095. DOI: [10.1021/ic00092a013](https://doi.org/10.1021/ic00092a013).
- [9] Brandt, K.; Porwollik-Czomperlik, I.; Siwy, M.; Kupka, T.; Shaw, R. A.; Davies, D. B.; Hursthouse, M. B.; Sykara, G. D. Thermodynamic vs Supramolecular Effects in the Regiocontrol of the Formation of New Cyclophosphazene-Containing Chiral Ligands with 1,1-Binaphthyl Units: Spiro vs Ansa Substitution at the  $N_3P_3$  Ring. *J. Am. Chem. Soc.* **1997**, *119*, 12432–12440. DOI: [10.1021/ja972372m](https://doi.org/10.1021/ja972372m).
- [10] Davies, D. B.; Clayton, T. A.; Eaton, R. J.; Shaw, R. A.; Egan, A.; Hursthouse, M. B.; Sykara, G. D.; Porwollik-Czomperlik, I.; Siwy, M.; Brandt, K. Chiral Configurations of Cyclophosphazenes. *J. Am. Chem. Soc.* **2000**, *122*, 12447–12457. DOI: [10.1021/ja0028469](https://doi.org/10.1021/ja0028469).
- [11] Coles, S. J.; Davies, D. B.; Eaton, R. J.; Kiliç, A.; Shaw, R. A.; Çiftçi, Y. G. Structural and Stereogenic Properties of Spiro- and Ansa-Substituted 1,3-Propanedioxy Derivatives of a Spermine-Bridged Cyclophosphazene. *Polyhedron* **2006**, *25*, 953–962. DOI: [10.1016/j.poly.2005.10.023](https://doi.org/10.1016/j.poly.2005.10.023).
- [12] Bešli, S.; Coles, S. J.; Davies, D. B.; Erkovan, A. O.; Hursthouse, M. B.; Kiliç, A. Single -, Double- and Triple-Bridged Derivatives of Cyclophosphazenes with an Octafluorohexane-1,6-Diol. *Polyhedron* **2009**, *28*, 3593–3599. DOI: [10.1016/j.poly.2009.07.042](https://doi.org/10.1016/j.poly.2009.07.042).
- [13] Finchan, J. K.; Hursthouse, M. B.; Parkes, H. G.; Shaw (Nee Gozen), L. S.; Shaw, R. A. The Structures of  $N_3P_3(NH_2)_2Cl_4$ ,  $N_3P_3(NPPH_3)(NH_2)Cl_4$  and  $N_3P_3(NPPH_3)Cl_5$  and a Comparison with Other Phosphazenylophosphazenes. The Relationship of Conformation in the Solid State to Conformation in Solution. *Acta Crystallogr. B Struct. Sci.* **1986**, *B42*, 462–476. DOI: [10.1107/S0108768186097872](https://doi.org/10.1107/S0108768186097872).
- [14] Yıldırım, T.; Bilgin, K.; Yenilmez Çiftçi, G.; Tanrıverdi Eçik, E.; Şenkuytu, E.; Uludağ, Y.; Tomak, L.; Kiliç, A. Synthesis, Cytotoxicity and Apoptosis of Cyclophosphazene Compounds as anti-Cancer Agents. *Eur. J. Med. Chem.* **2012**, *52*, 213–220. DOI: [10.1016/j.ejmech.2012.03.018](https://doi.org/10.1016/j.ejmech.2012.03.018).
- [15] Bovin, J. O.; Galy, J.; Labarre, J. F.; Sournies, F. Cyclophosphazenes as Novel Potential Antitumor Agents: x-Ray Crystal Structure of the Octapyrrolidino Cyclophosphazene,  $N_4P_4(NC_4H_8)_8$ . *J. Mol. Struct.* **1978**, *49*, 421–423. DOI: [10.1016/0022-2860\(78\)87282-2](https://doi.org/10.1016/0022-2860(78)87282-2).
- [16] Zhao, Z.; Guo, Q.; Li, X.; Sun, J.; Nie, Z.; Luo, W. Preparation and Properties of Novel Poly(Phosphazene-Aryl Amide)s Containing Cyclophosphazene Structures. *J. Appl. Polym. Sci.* **2013**, *128*, 4368–4377.
- [17] (a) Bovin, O. J.; Labarre, J. F.; Galy, J. The Crystal Structure of a New Antitumouragent: 2,2,4,4,6,6,8,8-Octapyrrolidinylcyclophosphazene,  $N_4P_4(NC_4H_8)_8$ . *Acta Crystallogr. B Struct. Crystallogr. Cryst. Chem.* **1979**, *35*, 1182–1186. doi:[10.1107/S0567740879005847](https://doi.org/10.1107/S0567740879005847). (b) Sournies, F.; Labarre, J.-F.; Spreafico, F.; Filipeschi, S.; Quan Jin, X. Attempts at the Production of More Selective Antitumourals. *J. Mol. Struct.* **1986**, *147*, 161–173. DOI: [10.1016/0022-2860\(86\)87067-3](https://doi.org/10.1016/0022-2860(86)87067-3).
- [18] Wolf, H. K.; Raad, M.; Snel, C.; Steenbergen, M. J.; Mhams, F.; Storm, G.; Henink, W. E. Biodegradable Poly(2-Dimethylamino Ethylamino)Phosphazene for in Vivo Gene Delivery to Tumor Cells. Effect of Polymer Molecular Weight. *Pharm. Res.* **2007**, *24*, 1572–1580. DOI: [10.1007/s11095-007-9299-z](https://doi.org/10.1007/s11095-007-9299-z).
- [19] Sournies, F.; Guerch, G.; Labarre, J. F. On the Antitumor Effectiveness of Aziridinocyclophosphazenes: crystal and Molecular Structure of the Non-Active 2,2-Bis (1-Aziridinyl)-4,4,6,6-Tetrachlorocyclophosphazene, Gem- $N_3P_3Az_2Cl_4$ . *J. Mol. Struct.* **1990**, *238*, 383–390. DOI: [10.1016/0022-2860\(90\)85029-I](https://doi.org/10.1016/0022-2860(90)85029-I).
- [20] Huizen, A. A. Aziridinyl Cyclophosphazenes. Ph.D. thesis, University of Groningen, The Netherlands, **1984**.
- [21] Sun, J.; Wang, X.; Wu, D. Novel Spirocyclic Phosphazene-Based Epoxy Resin for Halogen-Free Fire Resistance: Synthesis, Curing Behaviors, and Flammability Characteristics. *ACS Appl. Mater. Interfaces* **2012**, *4*, 4047–4061. DOI: [10.1021/am300843c](https://doi.org/10.1021/am300843c).
- [22] Yıldız, M.; Yılmaz, B.; Dolger, B. Synthesis, Spectral Properties, and Antimicrobial Activity of 2-Arilamino-2,4,4,6,6-Pentachloro-1,3,5,2,4,6-Triazatriphosphines and Poly[Bis(4-Fluorophenylamino)Phosphazene]. *Russ. J. Gen. Chem.* **2007**, *12*, 2117–2122. DOI: [10.1134/S1070363207120079](https://doi.org/10.1134/S1070363207120079).
- [23] İltter, E. E.; Asmafiliz, N.; Kılıç, Z.; Açık, L.; Yavuz, M.; Bali, E. B.; Solak, A. O.; Büyükkaya, F.; Dal, H.; Hőkelek, T. Syntheses, Structural and Electrochemical Investigations, Biological Activities and DNA Interactions of New Spirocyclic Monoferrocenylcyclophosphazenes. *Polyhedron* **2010**, *29*, 2933–2944. DOI: [10.1016/j.poly.2010.07.017](https://doi.org/10.1016/j.poly.2010.07.017).
- [24] Inoue, K.; Yamauchi, T.; Itoh, T.; Ihara, E. Ionic Conductivity of Cross-Linked Polymethacrylate Derivatives/Cyclophosphazenes/Li<sup>+</sup> Salt Complexes. *J. Inorg. Organomet. Polym. Mater.* **2007**, *17*, 367–375. DOI: [10.1007/s10904-007-9126-3](https://doi.org/10.1007/s10904-007-9126-3).
- [25] Barbera, J.; Bardaji, M.; Jimenez, J.; Laguna, A.; Martinez, M. P.; Oriol, L.; Serrano, J. L.; Zaragoza, I. Columnar Mesomorphic Organizations in Cyclophosphazenes. *J. Am. Chem. Soc.* **2005**, *127*, 8994–9002. DOI: [10.1021/ja051042w](https://doi.org/10.1021/ja051042w).
- [26] Görgülü, A. O.; Koran, K.; Özen, F.; Tekin, S.; Sandal, S. Synthesis, Structural Characterization and anti-Carcinogenic Activity of New Cyclophosphazenes Containing Dioxybiphenyl and Chalcone Groups. *J. Mol. Struct.* **2015**, *1087*, 1–10. DOI: [10.1016/j.molstruc.2015.01.033](https://doi.org/10.1016/j.molstruc.2015.01.033).
- [27] Baek, H.; Cho, Y.; Lee, C. O.; Sohn, Y. S. Synthesis and Antitumor Activity of Cyclophosphazene-(Diamine) Platinum(II) Conjugates. *Anticancer. Drugs* **2000**, *11*, 715–725. DOI: [10.1097/00001813-200010000-00008](https://doi.org/10.1097/00001813-200010000-00008).
- [28] Yenilmez Çiftçi, G.; Şenkuytu, E.; Bulut, M.; Durmuş, M. Novel Coumarin Substituted Water Soluble Cyclophosphazenes as “Turn-Off” Type Fluorescence Chemosensors for Detection of Fe<sup>3+</sup> Ions in Aqueous Media. *J. Fluoresc.* **2015**, *25*, 1819–1830. DOI: [10.1007/s10895-015-1672-4](https://doi.org/10.1007/s10895-015-1672-4).
- [29] Ozay, H.; Ozay, O. Synthesis and Characterization of Drug Microspheres Containing Phosphazene for Biomedical Applications. *Colloids Surf. A Physicochem. Eng. Asp* **2014**, *450*, 99–105. DOI: [10.1016/j.colsurfa.2014.03.022](https://doi.org/10.1016/j.colsurfa.2014.03.022).
- [30] Erdener Çıralı, D.; Uyar, Z.; Koyuncu, İ.; Hacıoğlu, N. Synthesis, Characterization and Catalytic, Cytotoxic and

- Antimicrobial Activities of Two Novel Cyclotriphosphazene-Based Multisite Ligands and Their Ru (II) Complexes. *Appl. Organometal. Chem.* **2015**, *29*, 536–542. DOI: [10.1002/aoc.3328](https://doi.org/10.1002/aoc.3328).
- [31] Tümer, Y.; Koç, L. Y.; Asmafiliz, N.; Kılıç, Z.; Hökelek, T.; Soltanzade, H.; Açık, L.; Yola, M. L.; Solak, A. O. Syntheses and Structural Investigations, Antimicrobial and Cytotoxic Activities and DNA Interactions of Vanillinato-Substituted NN or NO Spirocyclic Monoferrocenyl Cyclotriphosphazenes. *J. Biol. Inorg. Chem.* **2015**, *20*, 165–178. DOI: [10.1007/s00775-014-1223-5](https://doi.org/10.1007/s00775-014-1223-5).
- [32] Başterzi, N. S.; Bilge Koçak, S.; Okumuş, A.; Kılıç, Z.; Hökelek, T.; Çelik, Ö.; Türk, M.; Koç, L. Y.; Açık, L.; Aydın, B.; Dal, H. Syntheses, Structural Characterization and Biological Activities of Spiro-Ansa-Spiro-Cyclotriphosphazenes. *New J. Chem.* **2015**, *39*, 8825–8839. DOI: [10.1039/C5NJ01530H](https://doi.org/10.1039/C5NJ01530H).
- [33] Çil, E.; Tanyıldızı, M. A.; Özen, F.; Boybay, M.; Arslan, M.; Görgülü, A. O. Synthesis, Characterization, and Biological-Pharmacological Evaluation of New Phosphazenes Bearing Dioxypiphenyl and Schiff Base Groups. *Arch. Pharm. Pharm. Med. Chem.* **2012**, *345*, 476–485. DOI: [10.1002/ardp.201100412](https://doi.org/10.1002/ardp.201100412).
- [34] Is, Iklan, M.; Asmafiliz, N.; Özalp, E. E.; İlter, E. E.; Kılıç, Z.; C, Os, Ut, Bnyemin.; Yes, İlot, Serkan.; Kılıç, Adem.; Öztürk, Ash.; Hökelek, Tuncer.; et al. Synthesis, Structural Investigations, Biological Activities, and DNA Interactions of New N/O Spirocyclic Phosphazene Derivatives. The NMR Behaviors of Chiral Phosphazenes with Stereogenic Centers upon the Addition of Chiral Solvating Agents. *Inorg. Chem.* **2010**, *49*, 7057–7071. DOI: [10.1021/ic100781v](https://doi.org/10.1021/ic100781v).
- [35] Asmafiliz, N.; Kılıç, Z.; Hayvalı, Z.; Açık, L.; Hökelek, T.; Dal, H.; Öner, Y. Syntheses, Structural Investigations, Biological Activities, and DNA Interactions of New N/O Spirocyclotriphosphazenes. *Spectrochim. Acta Part A* **2012**, *86*, 214–223. DOI: [10.1016/j.saa.2011.10.027](https://doi.org/10.1016/j.saa.2011.10.027).
- [36] Brandt, K.; Kruszynski, R.; Bartczak, T.; Porwolik-Czomperlik, I. Porwolik-Czomperlik, I. AIDS- Related Lymphoma Screen Results and Molecular Structure Determination of a New Crown Ether Bearing Aziridinylcyclophosphazene, Potentially Capable of Ion-Regulated DNA Cleavage Action. *Inorg. Chim. Acta* **2001**, *322*, 138–144. DOI: [10.1016/S0020-1693\(01\)00557-6](https://doi.org/10.1016/S0020-1693(01)00557-6).
- [37] Siwy, M.; Se, K. D.; Kaczmarczyk, B.; Jaroszewicz, I.; Nasulewicz, A.; Pelczyńska, M.; Nevozhay, D.; Opolski, A. Synthesis and in Vitro Antileukemic Activity of Some New 1,3-(Oxytetraethylenoxy)Cyclotriphosphazene Derivatives. *J. Med. Chem.* **2006**, *49*, 806–810. DOI: [10.1021/jm0490078](https://doi.org/10.1021/jm0490078).
- [38] Okumuş, A.; Kılıç, Z.; Hökelek, T.; Dal, H.; Açık, L.; Öner, Y.; Koç, L. Y. Phosphorus–Nitrogen Compounds Part 22. Syntheses, Structural Investigations, Biological Activities and DNA Interactions of New Mono and Bis (4-Fluorobenzyl) Spirocyclophosphazenes. *Polyhedron* **2011**, *30*, 2896–2907. DOI: [10.1016/j.poly.2011.08.035](https://doi.org/10.1016/j.poly.2011.08.035).
- [39] Tümer, Y.; Asmafiliz, N.; Kılıç, Z.; Hökelek, T.; Koç, L. Y.; Açık, L.; Yola, M. L.; Solak, A. O.; Öner, Y.; DüNDAR, D.; Yavuz, M. Phosphorus–Nitrogen Compounds: Part 28. Syntheses Structural Characterizations, Antimicrobial and Cytotoxic Activities, and DNA Interactions of New Phosphazenes Bearing Vanillinato and Pendant Ferrocenyl Groups. *J. Mol. Struct.* **2013**, *1049*, 112–124.
- [40] Asmafiliz, N.; Kılıç, Z.; Öztürk, A.; Süzen, Y.; Hökelek, T.; Açık, L.; Çelik, Z. B.; Koç, L. Y.; Yola, M. L.; Üstündağ, Z. Phosphorus–Nitrogen Compounds: Part 25. Syntheses, Spectroscopic, Structural and Electrochemical Investigations, Antimicrobial Activities, and DNA Interactions of Ferrocenyldiaminocyclotriphosphazenes. *Phosphorus Sulfur Silicon Rel. Elem.* **2013**, *188*, 1723–1742. DOI: [10.1080/10426507.2013.779273](https://doi.org/10.1080/10426507.2013.779273).
- [41] Okumuş, A.; Bilge, S.; Kılıç, Z.; Öztürk, A.; Hökelek, T.; Yılmaz, F. Phosphorus–Nitrogen Compounds. Part 20: Fully Substituted Spiro-Cyclotriphosphazenic Lariat (PNP-Pivot) Ether Derivatives. *Spectrochim. Acta Part A* **2010**, *76*, 401–409. DOI: [10.1016/j.saa.2010.04.007](https://doi.org/10.1016/j.saa.2010.04.007).
- [42] Okumuş, A.; Elmas, G.; Cemaloğlu, R.; Aydın, B.; Binici, A.; Şimşek, H.; Açık, L.; Türk, M.; Güzel, R.; Kılıç, Z.; Hökelek, T. Phosphorus–Nitrogen Compounds. Part 35. Syntheses, Spectroscopic and Electrochemical Properties, Antituberculosis, Antimicrobial and Cytotoxic Activities of Mono-Ferrocenyl-Spiro Cycloctetraphosphazenes. *New J. Chem.* **2016**, *40*, 5588–5603. DOI: [10.1039/C6NJ00204H](https://doi.org/10.1039/C6NJ00204H).
- [43] Konar, V.; Yılmaz, Ö.; Öztürk, A. İ.; Kirbağ, S.; Arslan, M. Antimicrobial and Biological Effects of Bomphos and Phomphos on Bacterial and Yeast Cells. *Bioorg. Chem.* **2000**, *28*, 214–225. DOI: [10.1006/bioo.2000.1173](https://doi.org/10.1006/bioo.2000.1173).
- [44] Allcock, H. R.; Levin, M. L.; Austin, P. E. Quaternized Cyclic and High Polymeric Phosphazenes and Their Interactions with Tetracyanoquinodimethane. *Inorg. Chem.* **1986**, *25*, 2281–2288.
- [45] Moriya, K.; Suzuki, T.; Yano, S.; Miyajima, S. <sup>31</sup>P and <sup>13</sup>C NMR Studies of a Liquid-Crystalline Cyclotriphosphazene Derivative: Orientational Characteristics and Contrasting Shielding Anisotropies for Inorganic and Organic Moieties. *J. Phys. Chem. B* **2001**, *105*, 7920–7927. DOI: [10.1021/jp004299j](https://doi.org/10.1021/jp004299j).
- [46] Davarcı, Derya.; Beşli, Serap.; Demirbas, Erhan. Synthesis of a Series of Triple-Bridged Cyclotriphosphazene Hexa-Alkoxy Derivatives and Investigation of Their Structural and Mesomorphic Properties. *Liq. Cryst.* **2013**, *40*, 624–631. DOI: [10.1080/02678292.2013.773093](https://doi.org/10.1080/02678292.2013.773093).
- [47] Choi, Ji-Ae.; Kang, Yongku.; Kim, Dong-Won. Lithium Polymer Cell Assembled by in Situ Chemical Cross-Linking of Ionic Liquid Electrolyte with Phosphazene-Based Cross-Linking Agent. *Electrochim. Acta* **2013**, *89*, 359–364. DOI: [10.1016/j.electacta.2012.11.083](https://doi.org/10.1016/j.electacta.2012.11.083).
- [48] Labarre, J. F.; Guerch, G.; Sournies, F.; Lahana, R.; Enjalbert, R.; Galy, J. An Answer to the Spiro versus Ansa Dilemma in Cyclophosphazenes: Part IV. Reaction of Spermidine and Spermine on N<sub>3</sub>P<sub>3</sub>Cl<sub>6</sub>. *J. Mol. Struct.* **1984**, *116*, 75–88. DOI: [10.1016/0022-2860\(84\)80185-4](https://doi.org/10.1016/0022-2860(84)80185-4).
- [49] Labarre, J. F. Advances in Inorganic Chemistry. *Topics Curr. Chem.* **1985**, *U9*, 173–230.
- [50] Al-Madfa, Homaid A.; Hursthouse, Michael B.; Parkes, Harold G.; Shaw, Leyl[acaron] S.; Shaw, Robert A. The Reactions of N<sub>3</sub>P<sub>3</sub>L<sub>6</sub> with Polyfunctional Reagents. *Phosphorus Sulfur Silicon Rel. Elem.* **1986**, *28*, 203–211. DOI: [10.1080/03086648608072810](https://doi.org/10.1080/03086648608072810).
- [51] Al-Madfa, H. A.; Shaw, R. A. The Reactions of Hexachlorocyclotriphosphazatriene and Octachlorocycloctetraphosphazetate with Glycerol. *Phosphorus Sulfur Silicon Rel. Elem.* **1989**, *45*, 1–6. DOI: [10.1080/10426508908046069](https://doi.org/10.1080/10426508908046069).
- [52] Biddlestone, M.; Keat, R.; Parkes, H. G.; Rose, H.; Rycroft, D. S.; Shaw, R. A. Phosphorus–Nitrogen Compounds. Part 47. Conformations and Phosphorus–Phosphorus Spin–Spin Coupling Constants in Phosphazenylicyclophosphazenes. *Phosphorus Sulfur Silicon Rel. Elem.* **1985**, *25*, 25–31. DOI: [10.1080/03086648508074253](https://doi.org/10.1080/03086648508074253).
- [53] Krishnamurty, S. S.; Woods, M. *Nuclear Magnetic Resonance of Cyclophosphazenes (Annual Reports on NMR Spectroscopy)*; Academic Press Inc. Ltd.: London, **1987**; Vol. 19, pp 175–320. ISBN 0-12-505319-3.
- [54] Karplus, M. Contact Electron-Spin Coupling of Nuclear Magnetic Moments. *J. Chem. Phys.* **1959**, *30*, 11–15.
- [55] Karplus, M. J. Vicinal Proton Coupling in Nuclear Magnetic Resonance. *J. Am. Chem. Soc.* **1963**, *85*, 2870–2871. DOI: [10.1021/ja00901a059](https://doi.org/10.1021/ja00901a059).