



Biodiversity of the natural enemies of aphids (Hemiptera: Aphididae) in Northwest Turkey

Şahin Kök · Željko Tomanović ·
Zorica Nedeljković · Derya Şenal ·
İsmail Kasap

Received: 25 April 2019 / Accepted: 19 December 2019 / Published online: 2 January 2020
© Springer Nature B.V. 2020

Abstract In the present study, the natural enemies of aphids (Hemiptera: Aphididae) and their host plants including herbaceous plants, shrubs and trees were analysed to reveal their biodiversity and disclose tritrophic associations in different habitats of the South Marmara region of northwest Turkey. As a result of field surveys, 58 natural enemy species associated with 43 aphids on 58 different host plants were identified in the region between March of 2017 and November of 2018. In 173 tritrophic natural enemy-aphid-host plant interactions including association records new for Europe and Turkey, there were 21 representatives of the family Coccinellidae (Coleoptera), 14 of the family Syrphidae (Diptera) and 15 of the subfamily Aphidiinae

(Hymenoptera), as well as eight other generalist natural enemies. In these interactions, a total of 37 aphid-natural enemy associations—including 19 associations of *Acyrtosiphon pisum* (Harris) with natural enemies, 16 associations of *Therioaphis trifolii* (Monell) with natural enemies and two associations of *Aphis craccivora* Koch with natural enemies—were detected on *Medicago sativa* L. during the sampling period. Similarly, 12 associations of *Myzus cerasi* (Fabricius) with natural enemies were revealed on *Prunus avium* (L.), along with five associations of *Brevicoryne brassicae* (Linnaeus) with natural enemies (including mostly parasitoid individuals) on *Brassica oleracea* L. Also in the study, reduviids of the species *Zelus renardii* (Kolenati) are reported for the first time as new potential aphid biocontrol agents in Turkey. The results of field surveys show that the natural enemies of aphids have high biodiversity, which should be considered in the management of biological pest control.

Ş. Kök (✉) · İ. Kasap
Department of Plant Protection, Faculty of Agriculture, Çanakkale
Onsekiz Mart University, TR-17020 Çanakkale, Turkey
e-mail: sahinkok@gmail.com

e-mail: sahinkok@comu.edu.tr

Ž. Tomanović
Faculty of Biology, Institute of Zoology, University of Belgrade,
Studenstki trg 16, Belgrade 11000, Serbia

Z. Nedeljković
BioSense Institute-Research Institute for Information
Technologies in Biosystems, Dr Zorana Đinđića 1, Novi
Sad 21000, Serbia

D. Şenal
Department of Plant Protection, Faculty of Agriculture and Natural
Sciences, Bilecik Şeyh Edebali University,
TR-11230 Bilecik, Turkey

Keywords Biological control · Aphididae ·
Coccinellidae · Syrphinae · Aphidiinae · South Marmara

Introduction

As one of the most important of agricultural pests, aphids (Hemiptera, Aphididae) cause severe significant economic losses and are distributed in most terrestrial ecosystems throughout the world. They damage agricultural crops, forest plants and shade trees, especially in temperate regions, as a result of sap-sucking, honeydew

secretion and transmission of phytopathogenic viruses (Alford 2011; Diehl et al. 2013).

Natural enemies, including specialist and generalist species, play a very important role in suppressing populations of aphids and other arthropod herbivores (Symondson et al. 2002; Lu et al. 2012). Such natural enemies affect at least 50% of all aphid species on the surface of the earth (Waage 1991). Up to now, about 2000 arthropods, generally consisting of parasitoid species, have been determined as biological control agents worldwide (van Lenteren et al. 2006). Aphids are usually attacked by natural enemies such as parasitoids, predators and some pathogens. Of these, both the larvae and the adult stages of coccinellid species are predaceous on aphids, while only the larval stages of syrphids and chrysopids are predaceous on them. Also among parasitoids, species belonging to the family Braconidae (especially members of the subfamily Aphidiinae) develop as solitary endoparasitoids on aphid populations (Völkl et al. 2007).

Surveying the biodiversity of natural enemies of herbivore pests in both agricultural and natural habitats is quite important from the standpoint of planning biological control strategies. A richness of natural enemies makes possible stronger biological control in all habitats (Letourneau et al. 2009; Griffin et al. 2013; Katano et al. 2015). When the taxonomic distance between natural enemies on herbivore pests was high, the effect of this phenomenon was likewise positive (Griffin et al. 2013). However, greater diversity of natural enemies can lead to more ineffective pest control in agricultural habitats (Snyder and Wise 2001; Montoya et al. 2003).

In addition, the nature of associations of aphid and their host plants in different habitats can regulate the efficacy of influence exerted by specialist and generalist natural enemies on aphid populations. According to the hypothesis of host physiological phenotype, host plants that are nutrient-rich and short-lived tend to increase the density of aphids (Cronin et al. 2010), as in the case of populations of other nitrogen-limited herbivores (Mattson 1980), because host plants with a great amount of nitrogen have a positive effect on the development of aphid populations (Schütz et al. 2008). In view of the aforementioned considerations, the biodiversity of natural enemies and their interactions with both pests and host plants were here investigated in detail so as to better understand enemy-herbivore-host plant associations.

The aim of the present study was to reveal biodiversity of the natural enemies of pest aphids in different

habitats of the South Marmara region of northwest Turkey, which is an important crossroad between Europe and Asia and has a rich and diverse flora and fauna.

Material and methods

Sampling area

Our sampling area, the South Marmara region in northwest Turkey, consists of the Çanakkale and Balıkesir Provinces, which constitute an important link between Europe and Asia. This area includes 3% of Turkey's land, with a surface area of 783.562 km², and it is an outstanding region in terms of agricultural production. The region has a continental climate in the southeastern parts, milder climate in parts close to the coastal area and Mediterranean climate in regions of the gulf and islands. The sampling area also includes wilderness areas such as the Edremit Gulf, Biga Peninsula, Gallipoli Peninsula and Kaz Dağları (Ida Mountains), which are known to have a high degree of endemic floristic and faunistic diversity (Özhatay and Özhatay 2005).

Collection and identification of aphids and their natural enemies

Specimens of aphids and their natural enemies were collected from their host plants, including herbaceous plants, shrubs and trees in different habitats such as cultivated (simple landscapes-1), uncultivated (complex landscapes-2) and urban areas (3) in the Çanakkale and Balıkesir Provinces in northwest Turkey between March of 2017 and November of 2018. Accordingly, all samples were marked in the catalogue of associations. Sufficient numbers of apterous and alate specimens of aphids were transferred with a soft brush to Eppendorf tubes containing 70% ethyl alcohol for identification. The aphid specimens were identified by the first author using a LEICA DM 2500 microscope with a mounted HD camera and LAS software (version 4.1) according to Blackman and Eastop (2006, 2018) and Kök et al. (2016); Kök and Kasap (2019). Adults of natural enemies, especially ladybirds, hoverflies and generalist predators such as representatives of Hemiptera, Dermaptera and Neuroptera, were collected from host plants infested with aphids by visual and hand searching using sweep nets and suction tubes, after which they were brought to the laboratory in plastic

boxes and glass jars for preparation and identification. For sampling of predator larvae and parasitoids, different parts of aphid-infested plants including predator larvae and mummified aphids were brought in the same boxes to the laboratory for emergence of adult specimens. The larvae of predators were kept until they were adults in the aphid-containing plastic boxes, the upper part of which was covered with a net. For parasitoid emergence, mummified individuals in aphid colonies on the host plants were put in plastic bottles and kept in laboratory conditions (22.5 °C, 65% humidity, 16:8 L:D photoperiod). After the emergence of adult predators and parasitoids, the specimens of natural enemies of aphids were sent to competent experts for identification. Specimens belonging to the subfamily Aphidiinae (Braconidae) and families Syrphidae and Coccinellidae were identified by the second, third and fourth authors, respectively. Permanent slides of identified aphids and ladybird specimens were deposited in the library of slides maintained by the Department of Plant Protection, Faculty of Agriculture, Çanakkale Onsekiz Mart University. Specimens of other natural enemies were deposited in personal collections of the identifying experts.

Results and discussion

In determining biodiversity of the natural enemies of aphids in the present study, 58 natural enemy species associated with 43 aphids on 58 different host plants were recorded in different habitats of the South Marmara region of northwest Turkey during surveys conducted from 2017 to 2018. These enemies included 21 species of ladybird beetles of the family Coccinellidae (Coleoptera), whose members are one of the most frequently encountered natural enemies in all regions of Turkey. To be specific, Aslan and Uygün (2005) reported 33 coccinellid species feeding on 59 aphids in Kahramanmaraş and emphasized that the most common enemy is *Coccinella septempunctata* Linnaeus, which was found feeding on 41 aphid species. Also, Kök et al. (2017) determined that *C. septempunctata* is the most collected predator, with a 35,1% availability rate for aphids in Çanakkale, including our current survey area. These results are corroborated by the data we obtained in the present study, in which associations between *C. septempunctata* and 12 aphids were found in the South Marmara region. Species of the family Syrphidae, widely recognized as aphidophagous

hoverflies, are one of the most common known natural enemies of aphid populations (Chambers 1988) and are important agents for the biological control of many pests under both agricultural and greenhouse conditions (Pineda and García 2008; Leroy et al. 2010; Amorós-Jiménez et al. 2012). The most effective of them, larvae of the species *Episyrphus balteatus* de Geer, feed on about 100 aphid species throughout the world (Sadeghi and Gilbert 2000). In the same vein, Bayrak and Hayat (2008) stated that *E. balteatus* preys on 33 aphid species and is common in many regions of Turkey. Similarly, we found that the larvae of this hoverfly species feed on 10 aphid species harmful to many cultivated plants and are commonly encountered in our study area.

Among the most notable details in the catalogue of tritrophic relationships given below are the aphid-natural enemy associations on *Medicago sativa* L. Alfalfa (*M. sativa*) is the most widely used fodder crop worldwide (Walton 1983) and some aphid species such as *Acyrtosiphon pisum* (Harris); *Acyrtosiphon kondoi* Shinji; *Therioaphis trifolii* and *Aphis craccivora* have been reported as major pests of alfalfa fields in different parts of the world (Summers 1976; Nakashima and Akashi 2005; Rakhshani et al. 2006). These fields can also serve as rich reservoir areas for the natural enemies of aphids (Summers 1998) and researching the associations between aphids and their natural enemies in alfalfa fields is important to improve biological control strategies. As a result of sampling from different alfalfa fields, we detected a total of 37 associations of aphids with natural enemies, including 19 associations of *A. pisum* with natural enemies, 16 associations of *T. trifolii* with natural enemies and two associations of *A. craccivora* with natural enemies on *M. sativa* during the sampling period, as indicated in the catalogue below. Similarly, 12 associations of *M. cerasi* with natural enemies on *Prunus avium* (L.) were revealed in this study, along with five associations of *B. brassicae* with natural enemies (including mostly parasitoid individuals) on *Brassica oleracea* L.

The surprising finding of *Zelus renardii* (Kolenati), a generalist predator of leafhoppers, is one of the most interesting results of the present study. This generalist is a species native to North and Central America, from which it then began to spread to Hawaii and other tropical areas of the Pacific region (Weirauch et al. 2012). The predator later reached Greece and the Balkan Peninsula, which are close neighbours of Turkey (Davranoglou 2011; Simov et al. 2017). The first

finding of *Z. renardii* in the Turkish fauna was recorded by Çerçi and Koçak (2016) from İstanbul and İzmir, close to the European continent and the Balkan Peninsula. In another study, *Z. renardii* was determined as a natural enemy of *Glycaspis brimblecombei* (Moore) feeding on *Eucalyptus camaldulensis* Dehnh (Yurt and Karaca 2018). Finally, *Z. renardii* associated with *Cinara tujaefilina* (Del Guaricio) feeding on *Platycladus orientalis* is here determined for the first time as a predator of aphids in Turkey (in the South Marmara region). We feel that as a natural enemy, this generalist predator has the potential to be an agent for the biological control of aphids, and that records of its finding will be reported from different habitats and regions of Turkey in years to come. However, it should not be forgotten that *Z. renardii* is an intraguild predator of the larvae of *Chrysoperla carnea* (Stephens), an important agent for the biological control agent of aphids (Rosenheim et al. 1993; Cisneros and Rosenheim 1997).

On the other hand, parasitoid species, generally taxa belonging to the subfamily Aphidiinae (Hymenoptera: Braconidae), are specialist natural enemies of aphid populations and attack by laying their eggs in the body of nymphs and adult instars of their prey aphids (Bai and Mackauer 1990; Ralec 1995). Up to now, a number of authors have conducted various studies to determine the species of aphid parasitoids in many regions of Turkey, and they have contributed new records for the Turkish fauna (Ölmez and Ulusoy 2003; Aslan et al. 2004; Çetin Erdoğan et al. 2008; Tomanović et al. 2008; Kök et al. 2017). In the present study, we identified 15 parasitoid species from the subfamily Aphidiinae (Hymenoptera: Braconidae) associated with 26 aphids feeding on 30 host plant species. Also in this study, we collected as aphid natural enemies one species belonging to Chamaemyiida (Diptera) and two species of hyperparasitoids that cannot be identified. However, they were not included in the evaluation of tritrophic associations.

The interactions between aphids and their host plants are an important phenomenon affected by the nature of the habitat of host plants and aphids, location of the habitat and acceptance of aphids as prey by their natural enemies. The odours emitted by aphid-uninfested plants are a powerful guide enabling natural enemies to easily find their prey aphid on host plants in different habitats (Hatano et al. 2008). For example, allyl isothiocyanate and methyl salicylate emitted by the host plants *Brassica oleraceae* and *Glycine max* attract *Diaeretiella rapae*, *C. septempunctata* and syrphid flies, which are the

primary natural enemies of *B. brassicae* and *Aphis glycines* (Read et al. 1970; Zhu and Park 2005). Also, chemical and physical cues such as the alarm pheromone honeydew, its smell, aphid body colour and the shape and movement of aphids are effective factors enabling suitable prey aphids to be reached by both their parasitoids and their predators (Michaud and Mackauer 1994; Harmon et al. 1998). In light of the foregoing considerations, we have catalogued below the natural enemy-prey aphid-host plant associations in the South Marmara region of northwest Turkey so that these interactions in different habitats can be better understood and used more effectively in strategies for the biological control of aphids. In the present study, 173 tritrophic natural enemy-aphid-host plant associations (including association records new for Europe and Turkey) were determined that consisted of 58 natural enemies, 43 aphids and 58 different host plant species, as shown in the catalogue below.

Catalogue of plant-aphid-predator/parasitoid associations from the South Marmara region of northwest Turkey.

Amaranthus albus L. (Amaranthaceae)

Aphis craccivora Koch, 1854: *COCCINELLIDAE (Coleoptera): *Hippodamia variegata* Goeze, 1777 (1). *SYRPHIDAE (Diptera): *Paragus tibialis* (Fallén, 1817) (1). *Note: The association between *A. craccivora* and *P. tibialis* is recorded for the first time on host plants such as *A. albus*.

Amaranthus retroflexus L. (Amaranthaceae)

Aphis craccivora: *SYRPHIDAE (Diptera): *P. tibialis* (1). *Note: The association between *A. craccivora* and *P. tibialis* is recorded for the first time on host plants such as *A. retroflexus*.

Apiaceae (unidentified plant species)

Unidentified aphid species 1: *SYRPHIDAE (Diptera): *Episyrphus balteatus* (de Geer, 1776) (1).

Asteraceae (unidentified plant species)

Unidentified aphid species 2: *SYRPHIDAE (Diptera): *Eupeodes corollae* (Fabricius, 1794) (3); *Eupeodes*

luniger (Meigen, 1822) (2); *Meliscaeva auricollis* (Meigen, 1822) (3); *Scaeva pyrastris* (Linnaeus, 1758) (3).

Uroleucon sp.: *BRACONIDAE (Hymenoptera): *Aphidius funebris* Mackauer, 1961 (2); *Praon yomenae* Takada, 1968 (2).

Berberis sp. (Berberidaceae)

Liosomaphis berberidis (Kaltenbach, 1843): *COCCINELLIDAE (Coleoptera): *Coccinella septempunctata* Linnaeus, 1758 (3).

Berberis thunbergii DC. (Berberidaceae)

L. berberidis: *COCCINELLIDAE (Coleoptera): *C. septempunctata* (3); *Harmonia axyridis* Pallas, 1773 (3); *H. variegata* (3).

Brassica oleracea L. (Brassicaceae)

Brevicoryne brassicae (Linnaeus, 1758): *COCCINELLIDAE (Coleoptera): *C. septempunctata* (1). *SYRPHIDAE (Diptera): *E. corollae* (1); *S. pyrastris* (1); *Scaeva selenitica* (Meigen, 1822) (1). ***Note:** The association between *S. selenitica* and *B. brassicae* is recorded for the first time in Turkey. This relationship was previously documented only from Poland (Malinowska 1973; Rojo et al. 2003). *BRACONIDAE (Hymenoptera): *Diaeretiella rapae* (Curtis, 1860) (1).

Capsella rubella Reut. (Brassicaceae)

Aphis craccivora: *BRACONIDAE (Hymenoptera): *Lysiphlebus fabarum* (Marshall, 1896) (2); *Lysiphlebus testaceipes* (Cresson, 1880) (2).

Carlina sp. (Asteraceae)

Uroleucon jaceae (Linnaeus, 1758): *BRACONIDAE (Hymenoptera): *P. yomenae* (2).

Catalpa bignonioides Walter (Bignoniaceae)

Aphis catalpa Mamontova, 1953: *COCCINELLIDAE (Coleoptera): *Adalia bipunctata* (Linnaeus, 1758) (3).

Chenopodium album L. (Amaranthaceae)

Aphis fabae Scopoli, 1763: *BRACONIDAE (Hymenoptera): *Binodoxys angelicae* (Haliday, 1833) (1).

Aphis solanella Theobald, 1914: *COCCINELLIDAE (Coleoptera): *H. variegata* (2).

Hayhurstia atriplicis (Linnaeus, 1761): *SYRPHIDAE (Diptera): *E. balteatus* (1). *BRACONIDAE (Hymenoptera): *B. angelicae* (2); *D. rapae* (2).

Chrysanthemum sp. (Asteraceae)

Aphis gossypii Glover, 1877: *BRACONIDAE (Hymenoptera): *Aphidius colemani* Viereck, 1912 (3).

Crepis sp. (Asteraceae)

A. gossypii: *BRACONIDAE (Hymenoptera): *Aphidius matricariae* Haliday, 1834 (2).

Cupressus arizonica Greene (Cupressaceae)

Cinara fresai Blanchard, 1939: *COCCINELLIDAE (Coleoptera): *Adalia decempunctata* (Linnaeus, 1758) (3); *C. septempunctata* (3); *Nephus includens* (Kirsch, 1870) (3).

Cydonia oblonga mill. (Rosaceae)

Ovatus insitus (Walker, 1849): *COCCINELLIDAE (Coleoptera): *A. bipunctata* (1); *C. septempunctata* (1).

Cynara scolymus L. (Asteraceae)

Brachycaudus cardui (Linnaeus, 1758): *SYRPHIDAE (Diptera): *E. balteatus* (1).

Cynara sp. (Asteraceae)

B. cardui: *COCCINELLIDAE (Coleoptera): *C. septempunctata* (2). *SYRPHIDAE (Diptera): *S. selenitica* (2). ***Note:** This is a new aphid-syrphid association according to Rojo et al. 2003.

Cynoglossum creticum mill. (Boraginaceae)

Acyrtosiphon malvae (Mosley, 1841): *COCCINELLIDAE (Coleoptera): *C. septempunctata* (1); *Scymnus pallipediformis* Gunther, 1958 (1).

Euphorbia rigida M.Bieb. (Euphorbiaceae)

Aphis vallei Hille Ris Lambers & Stroyan, 1959: *COCCINELLIDAE (Coleoptera): *Scymnus apetzi* Mulsant, 1846 (2). *SYRPHIDAE (Diptera): *Scaeva albomaculata* (Macquart, 1842) (2). *Note: This is a new aphid-syrphid association. Additionally, this syrphid species is found for the first time on a host plant from the genus *Euphorbia* (Rojo et al. 2003). *BRACONIDAE (Hymenoptera): *Ephedrus persicae* Froggatt, 1904 (2).

Hedera helix L. (Araliaceae)

Aphis hederæ Kaltentbach, 1843: *BRACONIDAE (Hymenoptera): *B. angelicae* (3).

Hibiscus syriacus L. (Malvaceae)

A. gossypii: *BRACONIDAE (Hymenoptera): *L. testaceipes* (3).

Juniperus oxycedrus L. (Cupressaceae)

Cynara oxycedri Binazzi, 1996: *BRACONIDAE (Hymenoptera): *Pauesia* sp. Quilis, 1931 (3).

Lactuca sp. (Asteraceae)

Acyrtosiphon lactucae (Passerini, 1860): *BRACONIDAE (Hymenoptera): *A. matricariae* (2).

Malus domestica Borkh. (Rosaceae)

Dysaphis plantaginea (Passerini, 1860): *COCCINELLIDAE (Coleoptera): *A. bipunctata* (1); *N. includens* (1). *CHRYSOPIDAE (Neuroptera): *Chrysoperla carnea* (Stephens, 1836) (1). *BRACONIDAE (Hymenoptera): *A. matricariae* (1); *E. persicae* (1).

A. gossypii: *BRACONIDAE (Hymenoptera): *B. angelicae* (1); *L. testaceipes* (1).

Malus floribunda Siebold ex Van Houtte (Rosaceae)

Aphis pomi De Geer, 1773: *BRACONIDAE (Hymenoptera): *Lipolexis gracilis* Forster, 1862 (3); *Lysiphlebus cardui* (Marshall, 1896) (3); *L. fabarum* (3).

Malva sp. (Malvaceae)

Aphis umbrella (Börner, 1950): *BRACONIDAE (Hymenoptera): *A. matricariae* (2).

Malva sylvestris L. (Malvaceae)

A. umbrella: *BRACONIDAE (Hymenoptera): *Lysiphlebus confusus* Tremblay & Eady, 1978 (1).

Matricaria sp. (Asteraceae)

Unidentified aphid species 2: *BRACONIDAE (Hymenoptera): *L. testaceipes* (2).

Medicago sativa L. (Leguminosae)

Acyrtosiphon pisum (Harris, 1776): *COCCINELLIDAE (Coleoptera) (1): *A. decempunctata*; *C. septempunctata*; *Coccinula quatuordecimpustulata* (Linnaeus, 1758); *Exochomus nigromaculatus* Goeze, 1777; *Hippodamia tredecimpunctata* Linnaeus, 1758; *H. variegata*; *Propylea quatuordecimpunctata* (Linnaeus, 1758); *Psyllobora vigintiduopunctata* (Linnaeus, 1758); *S. apetzi*; *S. pallipediformis*; *Scymnus quadriguttatus* Fürsch et Kreissl, 1967. *SYRPHIDAE (Diptera) (1): *E. balteatus*; *E. corollae*; *Melanostoma mellinum* (Linnaeus, 1758); *S. pyrastris*; *Sphaerophoria scripta* (Linnaeus, 1758). *CHRYSOPIDAE (Neuroptera) (1): *C. carnea*. *BRACONIDAE (Hymenoptera) (1): *Aphidius ervi* Haliday, 1834; *Aphidius banksae* Kittel, 2016.

Aphis craccivora: *COCCINELLIDAE (Coleoptera): *S. apetzi* (1). *SYRPHIDAE (Diptera): *S. pyrastris* (1).

Therioaphis trifolii (Monell, 1882): *COCCINELLIDAE (Coleoptera) (1): *A. bipunctata*; *C. quatuordecimpustulata*; *E. nigromaculatus*; *H. variegata*; *P. quatuordecimpunctata*; *P. vigintiduopunctata*; *S. apetzi*; *S. pallipediformis*; *Scymnus rubromaculatus* (Goeze, 1778). *SYRPHIDAE (Diptera) (1): *E. corollae*; *S. scripta*; *Sphaerophoria rueppellii* Wiedemann, 1830. *Note: The association between *S. rueppellii* and *T. trifolii* is the first record in

Turkey. Before the present study, this association was confirmed only in Poland in Europe (Rojo et al. 2003). Also, the *E. corollae*-*T. trifolii* and *S. scripta*-*T. trifolii* associations are recorded for the first time in Turkey, having been previously known only from Poland in the whole Europe (Bankowska et al. 1975; Rojo et al. 2003). *MIRIDAE (Hemiptera) (1): *Deraeocoris serenus* (Douglas & Scott, 1868). *NABIDAE (Hemiptera): *Nabis pseudoferus* Remane, 1949. *CHRYSOPIDAE (Neuroptera) (1): *C. carnea*. *HEMEROBIIDAE (Neuroptera) (1).

Nerium oleander L. (Apocynaceae)

Aphis nerii Boyer de Fonscolombe, 1841: *COCCINELLIDAE (Coleoptera): *A. bipunctata* (3); *A. decempunctata* (3); *P. quatuordecimpunctata* (3). *BRACONIDAE (Hymenoptera): *A. colemani* (3); *B. angelicae* (3); *L. testaceipes* (3).

Aphis spiraeicola Patch, 1914: *COCCINELLIDAE (Coleoptera): *A. decempunctata* (3); *P. quatuordecimpunctata* (3).

Philadelphus coronarius L. (Hydrangeaceae)

Aphis fabae mordvilkoii Börner & Janich, 1922: *BRACONIDAE (Hymenoptera): *L. testaceipes* (3).

Pistacia terebinthus L. (Anacardiaceae)

Baizongia pistaciae (Linnaeus, 1767): *FORFICULIDAE (Dermaptera): *Forficula smyrnensis* Audinet-Serville, 1839 (1).

Platycladus orientalis (L.) Franco (Cupressaceae)

Cinara tujafilina (Del Guercio, 1909): *COCCINELLIDAE (Coleoptera): *A. decempunctata* (3); *C. septempunctata* (3); *H. axyridis* (3); *Harmonia quadripunctata* (Pontoppidan, 1763) (3); *Oenopia conglobata* (Linnaeus, 1758) (3); *S. pallipediformis* (3); *Scymnus subvillosus* (Goeze, 1777) (3). *ANTHOCORIDAE (Hemiptera): *Zelus renardii* (Kolenati, 1856) (3).

Portulaca oleracea L. (Portulacaceae)

Aphis craccivora: *SYRPHIDAE (Diptera): *P. tibialis* (1). *Note: The association between *A. craccivora* and

P. tibialis is recorded for the first time on host plants such as *P. oleracea*. *BRACONIDAE (Hymenoptera): *L. testaceipes* (1).

Prunus avium (L.) L. (Rosaceae)

Myzus cerasi (Fabricius, 1775): *COCCINELLIDAE (Coleoptera) (1): *A. bipunctata*; *Adalia fasciatopunctata revelieri* Mulsant, 1866; *A. decempunctata*; *Chilocorus bipustulatus* (Linnaeus, 1758); *H. axyridis*; *O. conglobata*. *SYRPHIDAE (Diptera) (1): *E. balteatus*; *Paragus pecchiolii* Rondani, 1857. *Note: The association of *M. cerasi* and *E. balteatus* is recorded for the first time in Turkey. Also, the association between *M. cerasi* and *P. pecchiolii* is recorded for the first time in Europe. *FORFICULIDAE (Dermaptera) (1): *Forficula aetolica* Brunner, 1882; *Forficula auricularia* Linnaeus, 1758; *F. smyrnensis*. *BRACONIDAE (Hymenoptera): *A. matricariae* (1).

Prunus domestica L. (Rosaceae)

Brachycaudus helichrysi (Kaltenbach, 1843): *COCCINELLIDAE (Coleoptera): *P. quatuordecimpunctata* (1). *SYRPHIDAE (Diptera): *E. balteatus* (1). *BRACONIDAE (Hymenoptera): *E. persicae* (1).

Hyalopterus pruni (Geoffroy, 1762): *COCCINELLIDAE (Coleoptera): *A. bipunctata* (1). *SYRPHIDAE (Diptera): *E. balteatus* (1).

Prunus dulcis (mill.) D.a. Webb (Rosaceae)

Hyalopterus amygdali (Blanchard, 1840): *COCCINELLIDAE (Coleoptera): *A. bipunctata* (3).

Prunus persica (L.) Batsch (Rosaceae)

Brachycaudus amygdalinus (Schouteden, 1905): *COCCINELLIDAE (Coleoptera): *C. septempunctata* (1).

Punica granatum L. (Lythraceae)

Aphis punicae Passerini, 1863: *COCCINELLIDAE (Coleoptera): *A. f. revelieri* (1). *BRACONIDAE (Hymenoptera): *A. matricariae* (1); *B. angelicae* (1); *L. testaceipes* (1).

Pyracantha coccinea M.Roem. (Rosaceae)

A. spiraecola: *COCCINELLIDAE (Coleoptera):
A. bipunctata (3); *H. axyridis* (3). *SYRPHIDAE (Dip-
 tera): *E. balteatus* (3). *Note: This is an aphid-syrphid
 association new in Turkey.

Rosa sp. (Rosaceae)

Macrosiphum euphorbiae (Thomas, 1878):
 *COCCINELLIDAE (Coleoptera): *C. septempunctata* (2).

Macrosiphum rosae (Linnaeus, 1758):
 *COCCINELLIDAE (Coleoptera): *C. septempunctata*
 (3). *SYRPHIDAE (Diptera): *E. balteatus* (2); *S.*
pyrastris (3). *Note: The association of *M. rosae* and
E. balteatus is recorded for the first time in Turkey.

Rumex crispus L. (Polygonaceae)

Aphis rumicis Linnaeus, 1758: *COCCINELLIDAE
 (Coleoptera): *Platynaspis luteorubra* (Goeze, 1777) (2).

Rumex pulcher L. (Polygonaceae)

A. rumicis: *BRACONIDAE (Hymenoptera):
B. angelicae (1); *L. fabarum* (1).

Rumex sp. (Polygonaceae)

A. solanella: *BRACONIDAE (Hymenoptera):
B. angelicae (2); *L. fabarum* (2).

Sambucus nigra L. (Adoxaceae)

Aphis sambuci Linnaeus, 1578: *COCCINELLIDAE
 (Coleoptera): *A. bipunctata* (3); *O. conglobata* (3).

Scabiosa sp. (Caprifoliaceae)

M. rosae: *COCCINELLIDAE (Coleoptera):
S. apetzi (2).

Silybum marianum (L.) Gaertn. (Asteraceae)

A. fabae: *BRACONIDAE (Hymenoptera):
L. testaceipes (2).

Solanum americanum Mill. (Solanaceae)

A. solanella: *SYRPHIDAE (Diptera): *S. pyrastris* (1).

Sonchus sp. (Asteraceae)

Hyperomyzus lactucae (Linnaeus, 1758):
 *SYRPHIDAE (Diptera): *E. balteatus* (1); *Paragus*
quadrifasciatus Meigen, 1822 (1). *Note: The associa-
 tion between *H. lactucae* and *P. quadrifasciatus* is re-
 corded for the first time in Turkey. This association was
 previously documented only in Spain (Marcos García
 1981, 1985; Rojo et al. 2003) and Portugal (Rojo et al.
 2003).

Spiraea vanhouttei (Briot) Zabel (Rosaceae)

A. spiraecola: *COCCINELLIDAE (Coleoptera):
A. bipunctata (3). *SYRPHIDAE (Diptera):
E. balteatus (3). *Note: This is an aphid-syrphid asso-
 ciation new in Turkey.

Tamarix sp. (Tamaricaceae)

Brachyunguis tamaricis (Lichtenstein, 1886):
 *COCCINELLIDAE (Coleoptera): *A. decempunctata*
 (3); *H. variegata* (3); *O. conglobata* (3).

Tribulus terrestris L. (Zygophyllaceae)

Aphis craccivora: *SYRPHIDAE (Diptera): *Paragus*
haemorrhous Meigen, 1822 (2). *Note: This is an
 aphid-syrphid relationship new in Turkey. The associa-
 tion was previously recorded only in Spain (Rojo and
 Marcos García 1998; Rojo et al. 2003) and Portugal
 (Gomes 1981; Rojo et al. 2003).

Triticum aestivum L. (Poaceae)

Sitobion avenae (Fabricius, 1775): *COCCINELLIDAE
 (Coleoptera): *C. quatuordecimpustulata* (1).

Urtica urens L. (Urticaceae)

A. solanella: *SYRPHIDAE (Diptera): *S. selenitica* (2).
 *Note: This is a new aphid-syrphid association accord-
 ing to Rojo et al. 2003.

Veronica sp. (Plantaginaceae)

A. gossypii: *COCCINELLIDAE (Coleoptera):
C. septempunctata (2). *BRACONIDAE (Hymenoptera):
B. angelicae (2).

Viburnum opulus L. (Adoxaceae)

A. spiraecola: *BRACONIDAE (Hymenoptera):
L. fabarum (3); *L. testaceipes* (3).

Viburnum tinus L. (Adoxaceae)

A. spiraecola: *COCCINELLIDAE (Coleoptera):
A. bipunctata (3).

Vicia faba L. (Leguminosae)

Aphis craccivora: *BRACONIDAE (Hymenoptera):
L. gracilis (1).

In conclusion, results of the present study reveal natural enemy-aphid-host plant associations and disclose biodiversity of the natural enemies of aphids in the South Marmara region of northwest Turkey, which is known for having a richly diversified fauna and flora. Fifty-eight natural enemies, 43 aphid species and 58 host plants were identified in the study, and 173 tritrophic associations of them were determined in all habitats. The vast majority of identified aphids and species of their natural enemies are new records for the fauna of the South Marmara region. We feel that determining biodiversity of the natural enemies of aphids—including harmful species that cause economically significant losses of crops—contributes both to more effective planning of strategies for biological control of these pests and to creation of an inventory of the region's faunal biodiversity. It is generally known that a high biodiversity of natural enemies strengthens the possibilities of achieving biological control. However, it should not be forgotten that the richness or absence of natural enemy biodiversity can have either a positive or a negative impact on the biological control of pests (Jonsson et al. 2017). Accordingly, it is believed that research in the future should be focused on the mechanisms governing interactions between natural enemies, pests and their host plants in order to increase the chances for success of biological pest control.

Acknowledgements This manuscript is a part of the PhD thesis of the first author. The authors are grateful to Associate Professor Dr. Sinan Anlaş and Dr. Gülten Yazıcı for identifying the generalist predator specimens; Associate Professor Dr. Ersin Karabacak for identifying the host plant species; and Mr. Raymond Dooley for proofreading the revised version of the manuscript and editing the English. Zorica Nedeljković's position at the University of Novi Sad is funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia (project OII 73002) and the H2020 project "ANTARES" (664387). The contribution of Željko Tomanović was supported by grant III43001 funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia.

Compliance with ethical standards The authors declare that the work is in compliance with ethical standards.

Conflict of interest The authors declare that the work involves no conflict of interests.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Alford, D. V. (2011). *Plant pests*. London: Harper Collins Publishers.
- Amarós-Jiménez, R., Pineda, A., Fereres, A., & Marcos-García, M. A. (2012). Prey availability and abiotic requirements of immature stages of the aphid predator *Sphaerophoria rueppellii*. *Biological Control*, 63, 17–24.
- Aslan, M. M., & Uygun, N. (2005). The aphidophagus coccinellid (Coleoptera: Coccinellidae) species in Kahramanmaraş, Turkey. *Turkish Journal of Zoology*, 29, 1–8.
- Aslan, M. M., Uygun, N., & Starý, P. (2004). A survey of aphid parasitoids in Kahramanmaraş, Turkey (Hymenoptera: Braconidae: Aphidiinae; and Hymenoptera: Aphelinidae). *Phytoparasitica*, 32(3), 255–263.
- Bai, B., & Mackauer, M. (1990). Oviposition and host-feeding patterns in *Aphelinus asychis* (Hymenoptera: Aphelinidae) at different aphid densities. *Ecological Entomology*, 15, 9–16.
- Bankowska, R., Kierych, E., Mikolajczyk, W., Palmowska, J., & Trojan, P. (1975). Aphid-aphidophage community in alfalfa cultures (*Medicago sativa* L.) in Poland. Part I. Structure and phenology of the community. *Annales Zoologici*, 32(14), 299–345.
- Bayrak, N., & Hayat, R. (2008). Faunistic studies on the species of Syrphidae (Diptera) in Kayseri province. *Bitki Koruma Bülteni*, 48(4), 35–49.
- Blackman, R. L., & Eastop, V. F. (2006). *Aphids on the world's herbaceous plants and shrubs*. Natural History Museum, London: John Wiley & Sons Ltd..
- Blackman R. L., & Eastop, V. F. (2018). *Aphids on the world's plants: an online identification and information guide*. <http://www.aphidsonworldsplants.info>

- Çerçi, B., & Koçak, Ö. (2016). Contribution to knowledge of the Heteroptera (Hemiptera) fauna of Turkey. *Journal of Insect Biodiversity*, 4(15), 1–18.
- Çetin Erdoğan, Ö., Tomanović, Ž., & Beyarslan, A. (2008). New aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) in the region of Marmara. *Acta Entomologica Serbica*, 13, 85–88.
- Chambers, R. J. (1988). Syrphidae. In A. K. Minks & P. Harrewijn (Eds.), *Aphids, their biology, natural enemies and control* (pp. 259–270). Amsterdam: Elsevier.
- Cisneros, J. J., & Rosenheim, J. A. (1997). Ontogenetic change of prey preference in the generalist predator *Zelus renardii* and its influence on predator-predator interactions. *Ecological Entomology*, 22, 399–407.
- Cronin, J. P., Welsh, M. E., Dekkers, M. G., Abercrombie, S. T., & Mitchel, C. E. (2010). Host physiological phenotype explains pathogen reservoir potential. *Ecology Letters*, 13, 1221–1232.
- Davranoglou, L. R. (2011). *Zelus renardii* (Kolenati, 1856), a new world reduviid discovered in Europe (Hemiptera: Reduviidae: Harpactorinae). *Entomologist's Monthly Magazine*, 147(1766–1768), 157–162.
- Diehl, E., Sereda, E., Wolters, V., & Birkhofer, K. (2013). Effects of predator specialization, host plant and climate on biological control of aphids by natural enemies: A meta-analysis. *Journal of Applied Ecology*, 50, 262–270.
- Gomes, A. (1981). Sirifeos colhidos em Portugal durante 1977 (Diptera: Syrphidae). *Agronomia Lusitana*, 41(1), 5–24.
- Griffin, J. N., Byrnes, J. E. K., & Cardinale, B. J. (2013). Effects of predator richness on prey suppression: A meta-analysis. *Ecology*, 94, 2180–2187.
- Harmon, J. P., Losey, J. E., & Ives, A. R. (1998). The role of vision and colour in the close proximity foraging behaviour of four coccinellid species. *Oecologia*, 115, 287–292.
- Hatano, E., Kunert, G., Michaud, J. P., & Weisser, W. W. (2008). Chemical cues mediating aphid location by natural enemies. *European Journal of Entomology*, 105, 797–806.
- Jonsson, M., Kaartinen, R., & Straub, C. S. (2017). Relationships between natural enemy diversity and biological control. *Current Opinion in Insect Science*, 20, 1–6.
- Katano, I., Doi, H., Eriksson, B. K., & Hillebrand, H. (2015). A cross-system meta analysis reveals coupled predation effects on prey biomass and diversity. *Oikos*, 124, 1427–1435.
- Kök, Ş., & Kasap, İ. (2019). Aphid (Hemiptera: Aphididae) species of the South Marmara region of Turkey including the first record of *Dysaphis radicola meridialis* Shaposhnikov, 1964 for the aphid fauna of Turkey. *Turkish Journal of Entomology*, 43(1), 63–78.
- Kök, Ş., Kasap, İ., & Özdemir, I. (2016). Aphid (Hemiptera: Aphididae) species determined in Çanakkale Province with a new record for the aphid fauna of Turkey. *Turkish Journal of Entomology*, 40(4), 397–412.
- Kök, Ş., Tomanović, Ž., Şenal, D., Baştuğ, G., & Kasap, İ. (2017). Aphidophagous coccinellid and parasitoid species determined in the Çanakkale Province with a new record for the parasitoid fauna of Turkey. *Bitki Koruma Bülteni*, 57(4), 485–502.
- Leroy, P. D., Verheggen, F. J., Capella, Q., Francis, F., & Haubruge, E. (2010). An introduction device for the aphidophagous hoverfly *Episyrphus balteatus* (De Geer) (Diptera: Syrphidae). *Biological Control*, 54, 181–188.
- Letourneau, D. K., Jedlicka, J. A., Bothwell, S. G., & Moreno, S. R. (2009). Effects of natural enemy biodiversity on the suppression of arthropod herbivores in terrestrial ecosystems. *Annual Review of Ecology, Evolution, and Systematics*, 40, 573–592.
- Lu, Y. H., Wu, K. M., Jiang, Y. Y., Guo, Y. Y., & Desneux, N. (2012). Widespread adoption of Bt cotton and insecticide decrease promotes biocontrol services. *Nature*, 487, 362–365.
- Malinowska, D. (1973). Larvae of hoverflies (Diptera: Syrphidae) in aphid colonies on some cultivated plants. *Polskie Pismo Entomologiczne*, 43, 607–619.
- Marcos García, M. A. (1981). Contribución al conocimiento de los Syrphidae (Diptera) de la zona noroeste de la provincia de Salamanca. *Bolletín de la Asociación española de Entomología*, 4, 157–171.
- Marcos García, M. A. (1985). Sirphidofauna de las sierras de Béjar, La Alberca y Gata. Subfamilia Syrphinae (Syrpinae, Diptera). *Revista Provincial de Estudios*, 16(17), 389–419.
- Mattson, W. J. (1980). Herbivory in relation to plant nitrogen content. *Annual Review of Ecology and Systematics*, 11, 119–161.
- Michaud, J. P., & Mackauer, M. (1994). The use of visual cues in host evaluation by aphidiid wasps. I. Comparison between 3 *Aphidius* parasitoids of the pea aphid. *Entomologia Experimentalis et Applicata*, 70, 273–283.
- Montoya, J. M., Rodriguez, M. A., & Hawkins, B. A. (2003). Food web complexity and higher-level ecosystem services. *Ecology Letters*, 6, 587–593.
- Nakashima, Y., & Akashi, M. (2005). Temporal and within-plant distribution of the parasitoid and predator complexes associated with *Acyrtosiphon pisum* and *A. kondoi* (Homoptera: Aphididae) on alfalfa in Japan. *Japanese Applied Entomology and Zoology*, 40(1), 137–144.
- Ölmez, S., & Ulusoy, M. R. (2003). A survey of aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) in Diyarbakır, Turkey. *Phytoparasitica*, 31(5), 524–528.
- Özhatay, N., & Özhatay, E. (2005). Kazdağı. In N. Özhatay, A. Byfield, & S. Atay (Eds.), *Türkiye'nin 122 Önemli Bitki Alanı* (pp. 73–76). İstanbul: Türkiye Doğal Hayatı Koruma Vakfı.
- Pineda, A., & García, M. (2008). Seasonal abundance of aphidophagous hoverflies (Diptera: Syrphidae) and their population levels in and outside Mediterranean sweet pepper greenhouses. *Annals of the Entomological Society of America*, 101(2), 384–391.
- Rakhshani, E., Talebi, A. A., Manzari, S., Rezwani, A., & Rakhshani, H. (2006). An investigation on alfalfa aphids and their parasitoids in different parts of Iran, with a key to the parasitoids (Hemiptera: Aphididae; Hymenoptera: Braconidae: Aphidiinae). *Journal of the Entomological Society of Iran*, 25(2), 1–14.
- Ralec, A. L. (1995). Egg contents in relation to host-feeding in some parasitic Hymenoptera. *Entomophaga*, 40, 87–93.
- Read, D. P., Feeny, P. P., & Root, R. B. (1970). Habitat selection by the aphid parasite *Diaretiella rapae* (Hymenoptera: Braconidae) and hyperparasite *Charips brassicae* (Hymenoptera: Cynipidae). *Canadian Entomologist*, 102, 1567–1578.
- Royo, S., & Marcos García, M. A. (1998). Catálogo de los sírfidos (Diptera: Syrphidae) afidófagos (Homoptera: Aphididae) presentes en los cultivos y plantas herbáceas de España y

- Portugal. *Bolletino di Zoologia agraria e di Bachicoltura, Ser. II*, 30(1), 39–54.
- Rojo, S., Gilbert, F. S., Marcos García, M. A., Nieto, J. M., & Mier, M. P. (2003). *A world review of predatory hoverflies (Diptera: Syrphidae: Syrphinae) and their prey* (319 pp). Alicante: Centro Iberoamericano de la Biodiversidad (CIBIO).
- Rosenheim, J. A., Wilhoit, L. R., & Armer, C. A. (1993). Influence of intraguild predation among generalist insect predators on the suppression of an herbivore population. *Oecologia*, 96, 439–449.
- Sadeghi, H., & Gilbert, F. (2000). Aphid suitability and its relationship to oviposition preference in predatory hoverflies. *Journal of Animal Ecology*, 69(5), 771–784.
- Schütz, K., Bonkowski, M., & Scheu, S. (2008). Effects of Collembola and fertilizers on plant performance (*Triticum aestivum*) and aphid reproduction (*Rhopalosiphum padi*). *Basic and Applied Ecology*, 9, 182–188.
- Simov, N., Gradinarov, D., & Davranoglou, L. R. (2017). Three new assassin bug records (Hemiptera: Heteroptera: Reduviidae) for the Balkan Peninsula. *Ecologica Montenegrina*, 13, 25–29.
- Snyder, W. E., & Wise, D. H. (2001). Contrasting trophic cascade generated by a community of generalist predators. *Ecology*, 82, 1571–1583.
- Summers, C. G. (1976). Population fluctuations of selected arthropods in alfalfa: Influence of two harvest practices. *Environmental Entomology*, 5(1), 103–110.
- Summers, C. G. (1998). Integrated pest management in forage alfalfa. *Integrated Pest Management Reviews*, 3, 127–154.
- Symondson, W. O. C., Sunderland, K. D., & Greenstone, M. H. (2002). Can generalist predators be effective biocontrol agents? *Annual Review of Entomology*, 47, 561–594.
- Tomanović, Ž., Beyarslan, A., Çetin Erdoğan, Ö., & Žikić, V. (2008). New records of aphid parasitoids (Hymenoptera, Braconidae, Aphidiinae) from Turkey. *Periodicum Biologorum*, 110(4), 335–338.
- van Lenteren, J. C., Bale, J., Bigler, F., Hokkanen, H. M. T., & Loomans, A. J. M. (2006). Assessing risks of releasing exotic biological control agents of arthropod pests. *Annual Review of Entomology*, 51, 609–634.
- Völkl, W., Mackauer, M., Pell, J. K., & Brodeur, J. (2007). Predators, parasitoids and pathogens. In H. F. van Emden & R. Harrington (Eds.), *Aphids as crop pests* (pp. 187–233). Oxford: Oxford University Press.
- Waage, J. K. (1991). Biodiversity as a resource for biological control. In D. L. Hawksworth (Ed.), *The biodiversity of microorganisms and invertebrates: Its role in sustainable agriculture* (pp. 149–163). Wallingford: CAB International.
- Walton, P. D. (1983). *Production and management of cultivated forages*. Reston: Reston Publishing Company, Inc..
- Weirauch, C., Alvarez, C., & Zhang, G. (2012). *Zelus renardii* and *Z. tetracanthus* (Hemiptera: Reduviidae): Biological attributes and the potential for dispersal in two assassin bug species. *Florida Entomologist*, 95(3), 641–649.
- Yurt, N. T., & Karaca, İ. (2018). İzmir ilinde Glycaspis brimblecombei (Moore) (Hemiptera: Aphalaridae)'nin Eucalyptus camaldulensis Dehnh üzerinde popülasyon değişimi ve bu alanda bulunan yararlı türler. *Süleyman Demirel Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 22, 83–88.
- Zhu, J. W., & Park, K. C. (2005). Methyl salicylate, a soybean aphid-induced plant volatile attractive to the predator *Coccinella septempunctata*. *Journal of Chemical Ecology*, 31, 1733–1746.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.