

## Aphidophagous coccinellid and parasitoid species determined in the Çanakkale Province with a new record for the parasitoid fauna of Turkey

Sahin KÖK<sup>1</sup>

Željko TOMANOVIĆ<sup>2</sup>

Derya ŞENAL<sup>3</sup>

Gökhan BAŞTUĞ<sup>1</sup> İsmail KASAP<sup>1</sup>

### ÖZ

#### Türkiye parazitoit faunası için yeni bir kayıt ile birlikte Çanakkale ilinde belirlenen afidofag coccinellid ve parazitoit türler

Bu çalışma 2013 ve 2015 yılları arasında Çanakkale ilinde afidofag coccinellid ve parazitoit türleri ile onların çeşitlilikleri ve birbirleri arasındaki tritrofik ilişkilerin (parazitoit/predatör-konukçu yaprakbiti-konukçu bitki) belirlenmesi amacıyla yürütülmüştür. Çalışma boyunca Aphidiinae (Hymenoptera: Braconidae) altfamilyasına ait altı cinsle bağlı 10 parazitoit tür ve Coccinellidae (Coleoptera) familyasına ait 10 cinsle bağlı 12 predatör tür tespit edilmiştir. Parazitoit türlerden *Aphidius hortensis* Marshall, 1896 Türkiye parazitoit faunası için yeni kayıttır. *Lysiphlebus testaceipes* (Cresson, 1880) (%32.9) ve *Lysiphlebus fabarum* (Marshall, 1896) (%23.0) türleri diğer parazitoit türlerden daha yoğun tespit edilmiştir. Benzer şekilde *Coccinella septempunctata* Linnaeus, 1758 (%35.1) ve *Adalia bipunctata* (Linnaeus, 1758) (%22.8) türleri predatör türler içerisinde en yaygın türler olarak belirlenmiştir. *Aphis fabae* Scopoli 1763 parazitoit türlerin konukçusu olarak en yaygın çeşitliliğe (Shannon-Wiener  $H = 0.658$ ) sahip iken *Cinara pini* (Linnaeus, 1758) ise predatör türlerin konukçusu olarak en yaygın çeşitliliğe sahiptir (Shannon-Wiener  $H = 1.386$ ). Bu çalışmada 14 farklı konukçu bitki üzerinde 18 parazitoit-yaprakbiti ilişkisi ve 19 farklı bitki üzerinde 28 predatör-yaprakbiti ilişkisi belirlenmiştir. *Berberis* sp. (Berberidaceae) üzerinde *A. hortensis* - *Liosomaphis berberidis* (Kaltenbach, 1843) ilişkisi Türkiye'de ilk kez bu çalışma ile kayıt edilmiştir.

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, Çanakkale Onsekiz Mart University, Çanakkale, Turkey

<sup>2</sup> Department of Invertebrate Zoology and Entomology, Faculty of Biology, University of Belgrade, Studentski trg 16, Belgrade, Serbia

<sup>3</sup> Department of Plant Protection, Faculty of Agriculture Sciences and Technology, Bilecik University, Turkey

Corresponding author e-mail: [sahinkok@gmail.com](mailto:sahinkok@gmail.com); [sahinkok@comu.edu.tr](mailto:sahinkok@comu.edu.tr)

Alınış (Received): 19.07.2017, Kabul ediliş (Accepted): 20.09.2017

**Anahtar kelimeler:** Yaprakbiti, coccinellid, parazitoit, *Aphidius hortensis*, Çanakkale

## ABSTRACT

This study was conducted between 2013 - 2015 to determine aphidophagous coccinellid and parasitoid species of aphids, their diversity and tritrophic associations (parasitoid/predator-host aphid-host plant) in the Çanakkale Province. In the course of the investigations, 10 species belonging to six genera of the subfamily Aphidiinae (Hymenoptera: Braconidae) were determined, in addition to 12 species belonging to 10 genera of the family Coccinellidae (Coleoptera). Among the parasitoids, *Aphidius hortensis* Marshall, 1896 is a new record for the Turkish parasitoid fauna. *Lysiphlebus testaceipes* (Cresson, 1880) (32.9%) and *Lysiphlebus fabarum* (Marshall, 1896) (23.0%) were more extensively found compared to other parasitoid species. Similarly, *Coccinella septempunctata* Linnaeus, 1758 (35.1%) and *Adalia bipunctata* (Linnaeus, 1758) (22.8%) were recorded as the most common predator species. *Aphis fabae* Scopoli 1763 had the greatest diversity among the parasitoids (Shannon-Wiener  $H = 0.658$ ), while *Cinara pini* (Linnaeus, 1758) had the greatest diversity among the predators (Shannon-Wiener  $H = 1.386$ ). As for tritrophic associations, 18 parasitoid-aphid associations on 14 host plants and 28 predator-aphid associations on 19 host plants were determined. The *A. hortensis* - *Liosomaphis berberidis* (Kaltenbach, 1843) association on *Berberis* sp. (Berberidaceae) is here newly recorded from Turkey. With the present study, the aphidophagous coccinellid and parasitoid species are determined for the first time in the Çanakkale Province of Turkey.

**Keywords:** Aphid, coccinellid, parasitoid, *Aphidius hortensis*, Çanakkale

## INTRODUCTION

Aphids (Hemiptera: Aphididae) are one of the most important pests causing damage to different plants in cultivated and uncultivated areas throughout the world. Aphids are responsible for direct damage to plants such as leaf curling, twisting of shoots, yellowing of leaves and stunting by sucking plant sap. In addition they also produce galls and deformities on plants by secreting toxic saliva during feeding. Moreover, they cause indirect damage by excreting honeydew on plant leaves and transmitting phytopathogenic viral diseases (Blackman and Eastop 2006, Jouraeva et al. 2006, van Emden and Harrington 2007). Recently, aphids have become more important in cultivated and uncultivated areas because they, winged aphids, migrate over greater distances and can produce several generations per annum (Gilbert et al. 2009, Kavallieratos et al. 2005, Posada-Flórez et al. 2014).

Aphid populations have a large number and various categories of specific and nonspecific natural enemies (parasitoids and predators) throughout the world. Among these natural enemies, parasitoid species belonging to the subfamilies Aphidiinae (Hymenoptera: Braconidae) and Aphelininae (Hymenoptera: Aphelinidae) are specialized to parasitize aphids and could play an important role in

biological control of aphids feeding on different plants (Aziza et al. 2014, Darsouei et al. 2011, Hagvar and Hofsvang 1991, Hughes 1989, Starý 2006, van Emden 1995, van Emden and Harrington 2007). Among aphid parasitoids, species belonging to the subfamily Aphidiinae are solitary wasp endoparasitoids of aphid populations, and they usually have a body length of 1-4 mm (Kavallieratos et al. 2001, Starý 1970, 1988). The subfamily Aphidiinae contains more than 400 species belonging to 60 genera and subgenera worldwide (Starý 1988). Many studies conducted in recent years, has led to the identification of a great number of new parasitoid species have in Southeast Europe, including Turkey (Akar and Çetin Erdoğan 2017, Çetin Erdoğan et al. 2008, 2010, Kavallieratos et al. 2003, Kavallieratos and Tomanović 2003, Tomanović and Starý 2001, Tomanović and Kavallieratos 2002, Tomanović et al. 2002, 2003a, 2003b). Predator species belonging to the family Coccinellidae (Coleoptera) are among the most important natural enemies of aphids and play a major role in control of aphid populations worldwide. Most of these species, generally known as ladybugs or ladybird beetles, feed as predators on both adult and immature stages of aphids and other pests. Also, some of these predators survive by phytophagy or mycethophagy (Chinery 1993, Tezcan and Uygun 2003). Predator species of the family Coccinellidae have been successfully used in biological control of many insect pests including aphids (Hemiptera: Aphididae), scale insects (Hemiptera: Coccoidea), whiteflies (Hemiptera: Aleyrodidae), thrips (Thysanoptera), mealybugs (Hemiptera: Pseudococcidae), leaf hoppers (Hemiptera: Cicadellidae) and other soft bodied insect pests worldwide (Magro et al. 2010).

Recently, synthetic pesticides have been heavily used in the control of aphid populations damaging different crops in Turkey and worldwide. These pesticides have caused serious harm to the environment (including beneficial insects) and human health (Desneux et al. 2004, 2006, 2007). Determining parasitoid and predator species as potential agents of biological control of aphid populations is therefore very important. The aim of the present study was to determine the species of parasitoids (Hymenoptera: Braconidae: Aphidiinae) and predator coccinellid (Coleoptera: Coccinellidae) of aphids feeding on different crops in cultivated and uncultivated areas of the Çanakkale Province, Turkey.

## MATERIALS AND METHODS

Aphid specimens, mummified aphids and predators were collected from herbaceous plants, shrubs and trees growing in cultivated and uncultivated areas from 2013 to 2015 in the Çanakkale Province (especially City Centre and Ayvacık, Bayramiç, Biga, Çan, Lapseki, Yenice), Turkey. Sampling was carried out aperiodically during spring and summer. The plant specimens colonized with aphids, mummified aphids and predators were brought to the laboratory in a plastic bag. Plants were pressed, herbarized and identified. Both apterous and alate aphids were collected from their host plants using a soft brush and put in an Eppendorf tube containing 70% alcohol to identify its species. Collection and preparation of aphids followed the method of

Hille Ris Lambers (1950), while identification of aphids was conducted according to Heie (1986), Bodenheimer and Swirski (1957) and Blackman and Eastop (2006, 2016). In the present study, the aphid species were determined by Kök et al. (2016).

For parasitoid emergence, mummified aphids and plant samples were put in plastic bottles inside a growth chamber (22.5 °C, 65% humidity, 16:8 L:D photoperiod). The lid of the plastic bottles had a circular opening covered with muslin cloth for ventilation purpose (Kavallieratos et al. 2004). After their emergence, the parasitoids were identified. Coccinellid species were collected from aphid-infested plant parts using a mouth aspirator in field conditions. Immature stages of the predators were brought to the laboratory and put in plastic bottles to complete their development. Preparation and identification of coccinellid species were done according to Uygun (1981).

All aphid, parasitoid and coccinellid specimens identified in the study were collected by the first author in the Çanakkale Province, Turkey. In the present study, the parasitoid and coccinellid species were identified by second and third author, respectively. Specimens of aphids and coccinellids were deposited in the insect collection of the Department of Plant Protection, Agricultural Faculty, Çanakkale Onsekiz Mart University, Turkey. Parasitoid specimens were deposited in that of the Institute of Zoology, Faculty of Biology, University of Belgrade, Belgrade, Serbia. Coordinates where we collected parasitoid and coccinellid species were recorded by GPS (Magellan Triton 500). The Shannon-Wiener  $H$  index was used to determine diversity of parasitoids and coccinellids of each aphid species in the Çanakkale Province. The nomenclature of diagnosed parasitoid species new for Turkey follows Sharkey and Wharton (1997).

## RESULTS AND DISCUSSION

This study was conducted between 2013 - 2015 to determine parasitoids and coccinellids of aphids, their diversity and tritrophic associations (parasitoid/coccinellid-host aphid-host plant) in the Çanakkale Province. During the study, 10 species belonging to six genera of the subfamily Aphidiinae (Hymenoptera: Braconidae) were determined, in addition to 12 species belonging to 10 genera of the family Coccinellidae (Coleoptera). Among the parasitoids, *Aphidius hortensis* Marshall, 1896 reared from *Liosomaphis berberidis* (Kaltenbach, 1843) feeding on *Berberis* sp. (Berberidaceae) is a new record for the Turkish parasitoid fauna. In the survey of parasitoid species, 18 associations of parasitoids-aphids occurring on 14 host plants were recorded. In the survey of coccinellid species, 28 associations of predators-aphids occurring on 19 host plants were reported. Moreover, the *A. hortensis* - *L. berberidis* association on *Berberis* sp. (Berberidaceae) was found as new record from Turkey. Specialized parasitoids of aphids of the genus *Liosomaphis* (Kavallieratos et al. 2004), the indicated species are of Palaearctic origin and were probably introduced into the USA and Canada (Smith 1944). It should be noted that parasitoids and predators of aphids are

determined for the first time in the Çanakkale Province. The taxonomy, host aphids, host plants and coordinates (Figure 1) are given below, along with the collection dates, the sex ratio and distribution of genera and species of parasitoids and coccinellids.




Figure 1. Location of sites where aphid parasitoids and coccinellids were sampled in the Çanakkale, Turkey.

### List of the species

#### Order Hymenoptera

#### Family Braconidae

#### Subfamily Aphidiinae

#### *Adialytus salicaphis* (Fitch, 1855)

Distribution: Andorra, Bulgaria, Canada, China, Croatia, Czech Republic, Finland, France, Georgia, Germany, Greece, Hungary, India, Iran, Iraq, Italy, Japan, Kazakhstan, Mexico, Moldova, Pakistan, Poland, Romania, Russia, Slovakia, Spain, Sweden, Tajikistan, Turkey, USA, United Kingdom, Uzbekistan, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Chaitophorus leucomelas* Koch 1854, *Populus* sp. (Salicaceae), 40°08'25.3" N 26°24'03.7" E, Çanakkale, City Centre, 10.VI.2015, 7♀♀, 2♂♂.

***Aphidius colemani* Viereck, 1912**

Distribution: Algeria, Angola, Argentina, Australia, Brazil, Bulgaria, Chile, China, Colombia, Costa Rika, Cyprus, Czech Republic, Egypt, France, Georgia, Germany, Greece, Hungary, India, Iran, Iraq, Israel, Italy, Japan, Jordan, Kenya, Lebanon, Libya, Madagascar, Morocco, Mozambique, Norway, Pakistan, Peru, Poland, Portugal, Russia, South Africa, Spain, Syria, Turkey, USA (Yu et al. 2012).

Material examined: *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°06'58.3" N 26°24'47.4" E, Çanakkale, City Centre, 12.VI.2015, 11♀♀, 2♂♂; *Aphis punicae* Passerini 1863, *Punica granatum* L. (Lythraceae), 40°08'24.7" N 26°24'10.8" E, Çanakkale, City Centre, 10.VI.2015, 4♀♀, 2♂♂.

**\**Aphidius hortensis* Marshall, 1896**

Distribution: Bulgaria, Canada, Czech Republic, Estonia, Finland, France, Georgia, Germany, Hungary, India, Italy, Moldova, the Netherlands, Poland, Slovakia, Spain, USA, Ukraine, United Kingdom, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Liosomaphis berberidis* (Kaltenbach 1843), *Berberis* sp. (Berberidaceae), 40°06'39.8" N 26°24'59.5" E, Çanakkale, City Centre, 27.VI.2015, 3♀♀.

*A. hortensis* is a new record for the aphid parasitoid fauna of Turkey. Also, the *A. hortensis* - *L. berberidis* association is a new record from Turkey. *A. hortensis* characterize short antennae with 14-15 antennal segments, elongated stigma, two-segmented labial palpomeres and short metacarpus (Figure 2). It is specialized parasitoids of *L. berberidis*.




Figure 2. *Aphidius hortensis* Marshall, 1896 (a) antennae, (b) head, frontal view, (c) ovipositor sheaths, lateral view, (d) petiole, dorsal view, (e) propodeum, dorsal view, (f) forewing.

***Aphidius matricariae* Haliday, 1834**

Distribution: Algeria, Andorra, Argentina, Belgium, Bulgaria, Canada, Chile, China, Cyprus, Czech Republic, Egypt, Finland, France, Germany, Hungary, India, Iran, Iraq, Israel, Italy, the Netherlands, Norway, Poland, Portugal, Serbia, Slovakia, Slovenia, Spain, Turkey, USA, Ukraine, United Kingdom, Yugoslavia (Yu et al. 2012).

Material examined: *Brachycaudus helichrysi* (Kaltenbach 1843), *Anthemis* sp. (Asteraceae), 40°08'36.5" N 26°25'01.4" E, Çanakkale, City Centre, 31.V.2015, 19♀♀, 20♂♂; *Brachycaudus helichrysi* (Kaltenbach 1843), *Helianthus annuus* L. (Asteraceae), 40°06'55.5" N 26°24'37.2" E, Çanakkale, City Centre, 15.V.2015, 3♀♀, 3♂♂.

***Aphidius rosae* Haliday, 1834**

Distribution: Andorra, Belarus, Belgium, Brazil, Bulgaria, Canada, China, Czech Republic, Finland, France, Germany, Greece, Hungary, India, Iran, Iraq, Ireland, Israel, Italy, the Netherlands, Norway, Pakistan, Poland, Portugal, Romania, Russia, Slovakia, Spain, Switzerland, Turkey, USA, Ukraine, United Kingdom, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Macrosiphum rosae* (Linnaeus 1758), *Rosa* sp. (Rosaceae), 40°05'59.6" N 26°22'03.3" E, Çanakkale, Kepez, 05.VI.2015, 23♀♀, 4♂♂.

***Binodoxys acalephae* (Marshall, 1896)**

Distribution: Andorra, Bulgaria, Canada, China, Czech Republic, Finland, France, Georgia, Germany, Hungary, India, Iran, Iraq, Italy, Kazakhstan, Moldova, Morocco, the Netherlands, Poland, Portugal, Russia, Serbia, Slovakia, Slovenia, Spain, Turkey, United Kingdom, Yugoslavia (Yu et al. 2012).

Material examined: *Aphis craccivora* Koch 1854, *Robinia pseudoacacia* L. (Fabaceae), 40°06'41.0" N 26°25'00.4" E, Çanakkale, City Centre, 06.VI.2015, ♀; *Aphis fabae* Scopoli 1763, *Robinia pseudoacacia* L. (Fabaceae), 40°06'41.0" N 26°25'00.4" E, Çanakkale, City Centre, 06.VI.2015, 5♀♀, 6♂♂.

***Lipolexis gracilis* Foerster, 1862**

Distribution: Algeria, Andorra, Bulgaria, China, Czech Republic, Finland, France, Georgia, Germany, Greece, Hungary, India, Italy, Japan, Kazakhstan, Lebanon, Moldova, the Netherlands, Pakistan, Poland, Portugal, Romania, Russia, Slovakia, Slovenia, Spain, Thailand, Turkey, Uzbekistan, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Aphis gossypii* Glover 1877, *Abelmoschus esculentus* (L.) Moench (Malvaceae), 40°13'56.1" N 26°32'25.4" E, Çanakkale, City Centre, 06.VI.2015, ♀.

***Lysiphlebus fabarum* (Marshall, 1896)**

Distribution: Afghanistan, Algeria, Andorra, Australia, Azerbaijan, Belgium, Bulgaria, China, Czech Republic, Egypt, Finland, France, Georgia, Germany, Greece, Hungary, India, Iran, Iraq, Israel, Italy, Japan, Kazakhstan, Korea, Lebanon, Moldova, Morocco, the Netherlands, Pakistan, Poland, Portugal, Romania, Russia, Slovakia, Slovenia, Spain, Switzerland, Syria, Turkey, Ukraine, United Kingdom, Uzbekistan, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Aphis fabae* Scopoli 1763, *Phaseolus vulgaris* L. (Fabaceae), 40°05'57.6" N 26°22'03.1" E, Çanakkale, Kepez, 05.VI.2015, 43♀♀; *Aphis fabae* Scopoli 1763, *Spartium junceum* L. (Fabaceae), 40°06'43.1" N 26°25'16.0" E, Çanakkale, University Campus, 23.V.2014, 4♀♀; *Aphis umbrella* (Börner 1950), *Malva* sp. (Malvaceae), 40°05'35.7" N 26°23'25.3" E, Çanakkale, Kepez, 25.IV.2014, 10♀♀; *Aphis gossypii* Glover 1877, *Abelmoschus esculentus* (L.) Moench (Malvaceae), 40°13'56.1" N 26°32'25.4" E, Çanakkale, Yapıldak, 06.VI.2015, 10♀♀.

***Lysiphlebus testaceipes* (Cresson, 1880)**

Distribution: Algeria, Argentina, Australia, Belgium, Brazil, Canada, Chile, China, Costa Rica, Croatia, Cuba, Czech Republic, France, Greece, India, Iran, Italy, Korea, Mexico, Pakistan, Peru, Portugal, South Africa, Spain, Turkey, USA, Uzbekistan, Yugoslavia, Serbia (Yu et al. 2012).

Material examined: *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°08'56.4" N 26°24'17.9" E, Çanakkale, City Centre, 11.VII.2014, 8♀♀, ♂; *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°06'59.3" N 26°24'49.0" E, Çanakkale, University Campus, 12.VI.2015, 13♀♀, 11♂♂; *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°10'00.1" N 26°24'40.4" E, Çanakkale, Esenler, 17.VI.2015, 2♀♀, ♂; *Aphis fabae* Scopoli 1763, *Vicia faba* L. (Fabaceae), 40°06'44.7" N 26°24'22.9" E, Çanakkale, Kepez, 14.V.2014, ♀, 2♂♂; *Aphis punicae* Passerini 1863, *Punica granatum* L. (Lythraceae), 40°05'55.5" N 26°22'04.1" E, Çanakkale, Kepez, 05.VI.2015, 10♀♀, 7♂♂; *Aphis punicae* Passerini 1863, *Punica granatum* L. (Lythraceae), 40°08'24.9" N 26°24'12.8" E, Çanakkale, City Centre, 10.VI.2015, 3♀♀, 6♂♂; *Aphis punicae* Passerini 1863, *Punica granatum* L. (Lythraceae), 40°09'03.0" N 26°24'29.6" E, Çanakkale, City Centre, 01.VI.2014, 14♀♀, 17♂♂.

***Pausia* sp. Quilis Perez, 1931**

Distribution: Afrotropical, East and West Palaearctic, European, Nearctic, Neotropical and Oriental regions (Yu et al. 2012).

Material examined: *Cinara pini* (Linnaeus 1758), *Pinus* sp. (Pinaceae), 40°06'37.7" N 26°24'37.1" E, Çanakkale, Kepez, 14.V.2014, 11♀♀, ♂.

## Order Coleoptera

### Family Coccinellidae

#### Subfamily Coccinellinae

##### *Adalia bipunctata* (Linnaeus, 1758)

Distribution: Middle Africa (Bielawski 1975), USA (Gordon 1985), Canada (Larson 2013), Palaearctic region, North Africa, North Asia and North China (Uygun 1981, Kovar 2007).

Material examined: *Aphis punicae* Passerini 1863, *Punica granatum* L. (Lythraceae), 40°13'56.1" N 26°32'25.4" E, Çanakkale, Yapıldak, 06.VI.2015, (2); *Brachycaudus helichrysi* (Kaltenbach), *Anthemis* sp. (Asteraceae), 40°08'36.5" N 26°25'01.4" E, Çanakkale, City Centre, 31.V.2015, (2); *Chaitophorus leucomelas* Koch, *Populus* sp. (Salicaceae), 40°06'32.9" N 26°24'59.3" E, Çanakkale, University Campus, 11.IV.2013, (4); *Cavariella aegopodii* (Scopoli), *Salix* sp. (Salicaceae), 40°06'48.9" N 26°25'14.2" E, Çanakkale, University Campus, 24.IV.2015, (2); *Cinara pini* (Linnaeus 1758), *Pinus* sp. (Pinaceae), 40°06'37.7" N 26°24'37.1" E, Çanakkale, Kepez, 14.V.2014 (1); *Hyalopterus pruni* (Geoffroy 1762), *Prunus persica* L. (Batsch) (Rosaceae), 40°01'38.1" N 27°02'54.2" E, Çanakkale, Çan, 07.VI.2013, (2).

##### *Adalia fasciatopunctata revelieri* Mulsant, 1866

Distribution: South Palaearctic region (Horion 1961).

Material examined: *Chaitophorus leucomelas* Koch, *Populus* sp. (Salicaceae), 40°06'32.9" N 26°24'59.3" E, Çanakkale, University Campus, 11.IV.2013, (3); *Cinara pini* (Linnaeus 1758), *Pinus* sp. (Pinaceae), 40°06'37.7" N 26°24'37.1" E, Çanakkale, Kepez, 14.V.2014, (1); *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°08'56.4" N 26°24'17.9" E, Çanakkale, City Centre, 11.VII.2014, (1); *Hyalopterus pruni* (Geoffroy 1762), *Prunus persica* L. (Batsch) (Rosaceae), 40°01'38.1" N 27°02'54.2" E, Çanakkale, Çan, 07.VI.2013, (1).

##### *Chilocorus bipustulatus* (Linnaeus, 1758)

Distribution: Mongolia (Bielawski 1975), USA (Gordon 1985), Palaearctic (especially Mediterranean region) (Uygun 1981, Kovar 2007), Sudan (Yousof et al. 2013),

Material examined: *Macrosiphum rosae* (Linnaeus 1758), *Rosa* sp. (Rosaceae), 40°08'44.2" N 26°24'59.6" E, Çanakkale, City Centre, 21.IV.2013, (5).

##### *Coccinella septempunctata* Linnaeus, 1758

Distribution: USA (Gordon 1985), Canada (Larson 2013), Sri Lanka (Ashfaquae 2012), Pakistan (Poorani 2004); Palaearctic and Afrotropical region (Horion 1961, Kovar 2007).

Material examined: *Aphis rumicis* Linnaeus 1758, *Rumex* sp. (Polygonaceae), 40°15'04.9" N 26°35'49.5" E, Çanakkale, Umurbey, 31.III.2013, (4); *Brevicoryne brassicae* (Linnaeus 1758), *Brassica oleracea* L. (Brassicaceae), 40°20'12.2" N 26°42'28.3" E, Lapseki, Beyçayırı, 31.III.2013, (4); *Aphis fabae* Scopoli 1763, *Vicia faba* L. (Fabaceae), 40°05'27.1" N 26°23'29.7" E, Çanakkale, Kepez, 14.III.2013, (5); *Hyperomyzus lactucae* (Linnaeus 1758), *Sonchus* sp. (Asteraceae), 40°6'40.6" N 26°24'53.1" E, Çanakkale, University Campus, 11.IV.2013, (1); *Aphis fabae solanella* Theobald 1914, *Solanum nigrum* L. (Solanaceae), 39°34'56.7" N 26°25'0.2" E, Çanakkale, Ayvacık, 10.IX.2013, (1); *Acyrtosiphon gossypii* Mordvilko 1914, *Sonchus asper* (L.) Hill 1769 (Asteraceae), 40°16'21.8" N 27°16'07.2" E, Çanakkale, Biga, 02.IV.2013, (5).

***Coccinella quatuordecimpustulata* (Linnaeus, 1758)**

Distribution: South Palaearctic region (Horion 1961).

Material examined: *Aphis fabae* Scopoli 1763, *Artemisia* sp. (Asteraceae), 40°4'46.30" N 26°21'41.6" E, Çanakkale, Dardanos campus, 18.IV.2013, (2).

***Exochomus (Parexochomus) nigromaculatus* Goeze, 1777**

Distribution: USA (Gordon 1985), Palaearctic region (Uygun 1981, Kovar 2007).

Material examined: *Hyperomyzus lactucae* (Linnaeus 1758), *Sonchus* sp. (Asteraceae), 40°6'40.6" N 26°24'53.1" E, Çanakkale, University Campus, 11.IV.2013, (1).

***Hippodamia variegata* Goeze, 1777**

Distribution: Palaearctic (especially western) region, Middle and North Africa, Europe, Arabia, India and China (Horion 1961, Kovar 2007).

Material examined: *Liosomaphis berberidis* (Kaltenbach), *Berberis* sp. (Berberidaceae), 40°06'39.8" N 26°24'59.5" E, Çanakkale, University Campus, 27.VI.2015, (2); *Aphis nerii* Boyer de Fonscolombe 1841, *Nerium* sp. (Apocynaceae), 40°08'56.4" N 26°24'17.9" E, Çanakkale, City Centre, 11.VII.2014, (1).

***Nephus (Bipunctatus) nigricans* Weise, 1879**

Distribution: Albania, France, Greece, Italy (Kovar 2007), Mediterranean region (Fursch 1965).

Material examined: *Aphis pomi* De Geer 1773, *Malus pumila* Miller, 1768 (Rosaceae), 39°46'7.9" N 26°46'45.7" E, Çanakkale, Bayramiç, 25.IV.2013, (1).

***Oenopia conglobata* (Linnaeus, 1758)**

Distribution: North America (Ashfaquae 2012), Europe, North Africa, Anatolia, Syria, Middle Asia and Siberia (Horion 1961).

Material examined: *Cinara pini* (Linnaeus 1758), *Pinus* sp. (Pinaceae), 40°06'37.7" N 26°24'37.1" E, Çanakkale, Kepez, 14.V.2014, (1); *Eucallipterus tiliae* (Linnaeus), *Tilia* sp. (Malvaceae), 40°06'36.4" N 26°24'49.5" E, Çanakkale, University Campus, 06.VI.2015, (1).

***Psyllobora vigintiduopunctata* (Linnaeus, 1758)**

Distribution: South Palaearctic region, Europe, North Africa, Anatolia, Armenia and Siberia (Horion 1961, Kovar 2007).

Material examined: *Myzus (Nectarosiphon) persicae* Sulzer 1776, *Capsicum* sp. (Solanaceae), 39°58'00.3" N 27°11'06.5" E, Çanakkale, Yenice, 04.IX.2013, (1).

***Scymnus pallipediformis* Gunther, 1958**

Distribution: Palaearctic region (Audisio and Canepari 2016).

Material examined: *Tinocallis (Sarucallis) kahawaluokalani* (Kirkaldy), *Lagerstroemia indica* L. Pers. (Lythraceae), 40°06'37.1" N 26°24'47.1" E, Çanakkale, University Campus, 27.VI.2015, (1).

***Scymnus (Pullus) subvillosus* (Goeze, 1777)**

Distribution: East Palaearctic, North Africa, Turkey, Russia, Georgia, Armenia, Azerbaijan, Syria, Israel, Iran, Iraq, Jordan, Arabian Peninsula (Audisio and Canepari 2016).

Material examined: *Cinara pini* (Linnaeus 1758), *Pinus* sp. (Pinaceae), 40°06'37.7" N 26°24'37.1" E, Çanakkale, Kepez, 14.V.2014, (1); *Chaitophorus leucomelas* Koch, *Populus* sp. (Salicaceae), 40°06'32.9" N 26°24'59.3" E, Çanakkale, University Campus, 11.IV.2013, (1).

In the present study, *Lysiphlebus testaceipes* (Cresson, 1880) (32.9%) and *Lysiphlebus fabarum* (Marshall, 1896) (23.0%) were found more extensively than the other parasitoid species. Similarly, *Coccinella septempunctata* Linnaeus, 1758 (35.1%) and *Adalia bipunctata* (Linnaeus, 1758) (22.8%) are recorded as the most common predator species. Interestingly, we did not find the widespread invasive species *Harmonia axyridis* Pallas, 1773. The diversity of parasitoids and predators associated with different host aphids was determined in all areas of the Çanakkale Province. The study revealed that the *Aphis fabae* Scopoli 1763 had the greatest diversity of parasitoids (Shannon-Wiener  $H = 0.658$ ) whereas *Cinara pini* (Linnaeus, 1758) had the greatest diversity of predators (Shannon-Wiener  $H = 1.386$ ) (Table 1).

Aphidophagous coccinellid and parasitoid species determined in the Çanakkale Province with a new record for the parasitoid fauna of Turkey

Table 1. Diversity and occurrence of parasitoids and predators on their host aphids in the Çanakkale Province, Turkey

Host Aphid Species	Parasitoid Species											Predator Coccinellid Species													
	<i>A. salicaphis</i>	<i>A. colemani</i>	<i>A. hortensis</i>	<i>A. matricariae</i>	<i>A. rosae</i>	<i>B. aculephae</i>	<i>L. gracilis</i>	<i>L. fabarum</i>	<i>L. testaceipes</i>	<i>Pauesi</i> asp.	Shanon-Weiner H	<i>A. bipunctata</i>	<i>A. f. revelieri</i>	<i>C. bipustulatus</i>	<i>C. septempunctata</i>	<i>C. quatuordecimpustulata</i>	<i>E. nigromaculatus</i>	<i>H. variegata</i>	<i>N. nigricans</i>	<i>O. conglobata</i>	<i>P. vigintiduopunctata</i>	<i>S. pallipediformis</i>	<i>S. subvillosus</i>	Shanon-Weiner H	
<i>A. craccivora</i>	-	-	-	-	-	1	-	-	-	-	0.000	-	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>A. fabae</i>	-	-	-	-	-	11	-	47	3	-	0.658	-	-	-	5	2	-	-	-	-	-	-	-	-	0.598
<i>A. fabaesolanella</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	1	-	-	-	-	-	-	-	-	-	0.000
<i>A. gossypii</i>	-	-	-	-	-	-	1	10	-	-	0.305	-	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>Acyr. gossypii</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	5	-	-	-	-	-	-	-	-	-	0.000
<i>A. nerii</i>	-	13	-	-	-	-	-	-	36	-	0.579	-	1	-	-	-	-	1	-	-	-	-	-	-	0.693
<i>A. pomi</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	-	-	-	1	-	-	-	-	-	-	0.000
<i>A. punicae</i>	-	6	-	-	-	-	-	-	57	-	0.314	2	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>A. rumicis</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	4	-	-	-	-	-	-	-	-	-	0.000
<i>A. umbrella</i>	-	-	-	-	-	-	-	10	-	-	0.000	-	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>B. helichrysi</i>	-	-	-	45	-	-	-	-	-	-	0.000	2	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>B. brassicae</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	4	-	-	-	-	-	-	-	-	-	0.000
<i>C. aegopodii</i>	-	-	-	-	-	-	-	-	-	-	0.000	2	-	-	-	-	-	-	-	-	-	-	-	-	0.000
<i>C. leucomelas</i>	9	-	-	-	-	-	-	-	-	-	0.000	4	3	-	-	-	-	-	-	-	-	-	-	1	0.974
<i>C. pini</i>	-	-	-	-	-	-	-	-	12	-	0.000	1	1	-	-	-	-	-	-	1	-	-	1	1.386	
<i>E. tiliae</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	-	-	-	-	1	-	-	-	-	-	0.000
<i>H. pruni</i>	-	-	-	-	-	-	-	-	-	-	0.000	2	1	-	-	-	-	-	-	-	-	-	-	-	0.637
<i>H. lactucae</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	1	-	1	-	-	-	-	-	-	-	0.693
<i>L. berberidis</i>	-	-	3	-	-	-	-	-	-	-	0.000	-	-	-	-	-	-	2	-	-	-	-	-	-	0.000
<i>M. rosae</i>	-	-	-	-	27	-	-	-	-	-	0.000	-	-	5	-	-	-	-	-	-	-	-	-	-	0.000
<i>M. persicae</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	-	-	-	-	-	-	1	-	-	-	0.000
<i>T. kahawaluokalani</i>	-	-	-	-	-	-	-	-	-	-	0.000	-	-	-	-	-	-	-	-	-	-	-	1	-	0.000
Occurrence ratio (%)	3.0	6.5	1.0	15.5	9.3	4.1	0.3	23.0	32.9	4.1	22.8	10.5	8.8	35.1	3.5	1.8	5.3	1.8	3.5	1.8	1.8	3.5			
Total number of parasitoids: 291 specimens											Total number of predators: 57 specimens														

A great number of local studies have been conducted to identify aphid parasitoids and predators in cultivated and uncultivated areas of Turkey. Ölmez and Ulusoy (2003) reported that 16 parasitoid species belonging to nine genera of the subfamily Aphiidinae (Hymenoptera: Braconidae) were determined in the Diyarbakır Province. Among these parasitoid species, *Monoctonus mali* van Achterberg 1989 was a new record for the aphid parasitoid fauna of Turkey. Aslan et al. (2004) investigated aphid parasitoids in the Kahramanmaraş Province. They reported 19 aphid parasitoid species belonging to the subfamily Aphiidinae (Hymenoptera: Braconidae) and family Aphelinidae (Hymenoptera) on different plants. A list of the aphid hosts of parasitoid species was given in the indicated study, and the finding of *Pauesia picta* (Haliday, 1834) was a new record for the aphid parasitoid fauna of Turkey. Güz and Kılınçer (2005) reported 10 aphid parasitoid species belonging to six genera of the family Braconidae (Hymenoptera) family in the Ankara Province. Çetin Erdoğan et al. (2008) announced that 17 species of Aphiidinae parasitoids were identified from region of Marmara including Provinces of Adapazarı, Balıkesir, Bursa, Bilecik, Çanakkale, Edirne, İstanbul, İzmit, Tekirdağ and Yalova. Tomanović et al. (2008) conducted a study to identify aphid parasitoids in the Marmara and Black Sea regions of Turkey. In this study, they recorded eight species new for the aphid parasitoid fauna of Turkey. Akar and Çetin Erdoğan (2017) reported that 27 species belongs to Aphiidinae were determined in eight different habitats in the Province of Edirne. Among these, seven species were new records for the Turkish Aphiidinae fauna. Baştuğ and Kasap (2015) surveyed species of the family Coccinellidae (Coleoptera) on different plants in cultivated and uncultivated areas of the province investigated in our study (the Çanakkale Province). They reported 20 coccinellid species by focusing only on the existence of coccinellid species in the Çanakkale Province, but not investigated aphid-coccinellid associations. Aslan and Uygun (2005) identified 33 predator coccinellid species associated with 59 aphids feeding on different plants in cultivated and uncultivated areas of the Kahramanmaraş Province. Many studies have been carried out to identify species of aphid parasitoids and predators in different districts of Turkey (Aslan and Karaca 2005, Ayyıldız and Atlıhan 2006, Güncan et al. 2010, Şahbaz and Uysal 2006). Although many local areas in Turkey have been investigated to determine parasitoids and predators of aphids, the number of their natural enemies is still quite limited.

In view of the negative effects of pesticides used to control of aphids and other pests on the environment and human health, parasitoids and predators of these pests have become more important in biological control studies. Protecting and supporting the native parasitoids and predators of aphids will increase the success of biological control efforts in field and greenhouse conditions. In addition to determining natural enemies of aphids feeding on plants in agricultural fields, aphid-parasitoid-predator associations on plants in uncultivated areas should also be investigated. With the present study, parasitoids and predators of aphids are determined for the first time in cultivated and uncultivated areas of the Çanakkale Province in Turkey. It is recommended that local fauna studies of parasitoids and predators of aphids be

conducted so that these natural enemies can be used more effectively in biological control studies. More comprehensive studies will contribute to knowledge of the fauna of natural enemies of aphids in Turkey.

### ACKNOWLEDGEMENTS

Contribution of Željko TOMANOVIĆ was supported by the Serbian Ministry of Education, Science and Technological Development (grant III43001).

### REFERENCES

- Akar S. and Çetin Erdoğan Ö. 2017. Türkiye Aphidiinae Faunasına (Hymenoptera: Braconidae) Yeni Kayıtlar ile Katkılar. *Trakya University Journal of Natural Sciences*, 18 (2) (In-press).
- Ashfaq A. 2012. Taxonomic Studies of Family Coccinellidae (Coleoptera) of Gigit-Baltistan, Pakistan. Department of Plant Protection. Faculty of Crop Protection Sciences. The University of Agriculture, Peshawar. 196p.
- Aslan B. and Karaca I. 2005. Fruit Tree Aphids and Their Natural Enemies in Isparta Region, Turkey. *Journal of Pest Science*, 78, 227-229.
- Aslan M. M. and Uygun N. 2005. The Aphidophagus Coccinellid (Coleoptera: Coccinellidae) Species in Kahramanmaraş, Turkey. *Turkish Journal of Zoology*, 29, 1-8.
- Aslan M. M., Uygun N. and Starý P. 2004. A Survey of Aphid Parasitoids in Kahramanmaraş, Turkey (Hymenoptera: Braconidae: Aphidiinae; and Hymenoptera: Aphelinidae). *Phytoparasitica*, 32 (3), 255-263.
- Audisio P. and Canepari C. 2016. Fauna Europaea: Coleoptera: Coccinellidae. Fauna Europaea version 2.6, (Web page: <http://www.faunaeur.org>) (Date accessed: September 2016).
- Ayyıldız Y. and Atlıhan R. 2006. Balıkesir İli Sebze Alanlarında Görülen Yaprakbiti Türleri ve Doğal Düşmanları. *Yüzüncü Yıl Üniversitesi Ziraat Fakültesi Tarım Bilimleri Dergisi*, 16 (1), 1-5.
- Aziza M.El-G., Ahmed S.A., El-Bassioni M.N., Mahfouz H.M. and El-Deeb M.G.A. 2014. Survey of Aphid Species and Their Associated Parasitoids and Predators on Wild Plants in North Sinai Governorate, Egypt. *Egyptian Journal of Biological Pest Control*, 24 (2), 283-288.
- Baştuğ G. and Kasap İ. 2015. Çanakkale İli Coccinellidae (Coleoptera) Familyası Üzerine Faunistik Çalışmalar. *Türkiye Biyolojik Mücadele Dergisi*, 6 (1), 41-50.
- Bielawski R. 1975. Ergebnisse der Zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei Nr. 352. Coccinellidae V und VI. *Fragmenta Faunistica* 20 (16), 247-271.
- Blackman R.L. and Eastop V.F. 2006. *Aphids on the World's Herbaceous Plants and Shrubs*. Wiley and Sons Ltd., London, UK, 1439 pp.

- Blackman R.L. and Eastop V.F. 2016. Aphids on the World's Plants: An Online Identification and Information Guide, <http://www.aphidsonworldsplants.info> (Date accessed: June 2016).
- Bodenheimer F.S. and Swirski E. 1957. The Aphidoidea of the Middle East. Weizmann Science Press of Israel, Jerusalem, 378 pp.
- Chinery M. 1993, Insects of Britain and Western Europe. Collins, London, 320 pp.
- Çetin Erdoğan Ö., Tomanović Ž. and Beyarslan A. 2008. New Aphid Parasitoids (Hymenoptera: Braconidae: Aphidiinae) in the Region of Marmara. Acta Entomologica Serbica, 13, 85 - 88.
- Çetin Erdoğan Ö., Tomanović Ž. and Beyarslan A. 2010. New Distributional Records on the Subfamily Aphidiinae (Hymenoptera: Braconidae) in Black Sea Region, Turkey. Linzer Biologische Beiträge, 42, 613-616.
- Darsouei R., Karimi J. and Modarres-Awal M. 2011. Parasitic Wasps as Natural Enemies of Aphid Populations in the Mashhad Region of Iran: New Data from DNA Barcodes and SEM. Archives of Biological Sciences, 63 (4), 1225-1234.
- Desneux N., Decourtye A. and Delpuech J.M. 2007. The Sublethal Effects of Pesticides on Beneficial Arthropods. Annual Review of Entomology, 52, 81-106.
- Desneux N., Denoyelle R. and Kaiser L. 2006. A Multi-Step bioassay to Assess the Effect of Deltamethrin on the Parasitic Wasp *Aphidius ervi*. Chemosphere, 65, 1697-1706.
- Desneux N., Pham-Dele`gue M.H. and Kaiser L. 2004. Effects of Sublethal and Lethal Doses of Lambda-Cyhalothrin on Oviposition Experience and Host Searching Behaviour of a Parasitic Wasp *Aphidius ervi*. Pest Management Science, 60, 381-389.
- Fursch H. 1965. Die Palearktischen Arten der Scymnus-bipunctatus-Gruppe und die Europaeschen Vertreter der Untergattung Sidis (Col. Cocc.). Mitt. Münch. Ent. Gesell. 55, Jahrg. 178-213.
- Gilabert A., Simon J.C., Mieuze L., Halkett F., Stoeckel S., Plantegenest M. and Dedryver C.A. 2009. Climate and Agricultural Context Shape Reproductive Mode Variation in an Aphid Crop Pest. Molecular Ecology, 18 (14), 3050-3061.
- Gordon R.D. 1985. The Coccinellidae of America North of Mexico. Journal of the New York Entomological Society, 93 (1), 1-912.
- Günçan A., Yoldaş Z. and Madanlar N. 2010. İzmir'de Şeftali Bahçelerinde Bulunan Yaprakbiti (Hemiptera: Aphididae) Türleri ve Doğal Düşmanları Üzerinde Araştırmalar. Türkiye Entomoloji Dergisi, 34 (3), 399-408.
- Güz N. and Kılınçer N. 2005. Aphid Parasitoids (Hymenoptera: Braconidae: Aphidiinae) on Weeds from Ankara, Turkey. Phytoparasitica, 33 (4): 359-366.
- Hagvar E.B. and Hofsvang T. 1991. Aphid Parasitoids (Hymenoptera: Aphidiidae): Biology, Host Selection and Use in Biological Control. Biocontrol News and Information, 12, 13-41.

- Heie O.E. 1986. The Aphidoidea (Hemiptera) of Fennoscandia and Denmark (III), Family Aphididae: Subfamily Pterocommatinae and Tribe Aphidini of Subfamily Aphidinae. E.J. Brill/Scandinavian Science Press Ltd., Leiden-Copenhagen, 314 pp.
- Hille Ris Lambers D. 1950. On Mounting Aphids and Other Soft-Skinned Insects. Entomologische Berichten, XIII, 55-58.
- Horion A. 1961. Faunistif der Mitteleuropäischen Käfer. Band VIII. Überlingen-Bodensee, Kommissionsverlag Buchdruckerei Ang. Feysel., 283-365.
- Hughes R.D. 1989. Biological Control in the Open Field In: Minks A.K. and Harrewijn P. (eds.), Aphids, Their Biology, Natural Enemies and Control, Crop Pests, Vol. 2C, Elsevier, Amsterdam, 167-198 p.
- Jouraeva V.A., Johnson D.L., Hassett J.P., Nowak D.J., Shipunova N.A. and Barbarossa D. 2006. Role of Sooty Mold Fungi in Accumulation of Fine-Particle-Associated PAHs and Metals on Deciduous Leaves. Environmental Research, 102, 272-282.
- Kavallieratos N.G. and Tomanović Ž. 2003. Some Rare and Endemic Aphid Parasitoid Species (Hymenoptera: Braconidae: Aphidiinae) from the Balkan Peninsula. Acta Entomologica Serbica, 6, 121-129.
- Kavallieratos N.G., Athanassiou C.G. and Tomanović Ž. 2003. A New Species and a Key to Greek *Praon* Haliday (Hymenoptera: Braconidae: Aphidiinae). Deutsche Entomologische Zeitschrift, 50, 13-22.
- Kavallieratos N.G., Athanassiou C.G., Tomanović Ž., Sciarretta A., Trematerra P. and Žikić V. 2005. Seasonal Occurrence, Distribution and Sampling Indices for *Myzus persicae* (Hemiptera: Aphidoidea) and Its Parasitoids (Hymenoptera: Braconidae: Aphidiinae) on Tobacco. European Journal of Entomology, 102 (3), 459-68.
- Kavallieratos N.G., Lykouressis D.P., Sarlis G.P., Stathas G.J., Segovia A.S. and Athanassiou C.G. 2001. The Aphidiinae (Hymenoptera: Ichneumonoidea: Braconidae) of Greece. Phytoparasitica, 29, 306-340.
- Kavallieratos N.G., Tomanović Ž., Starý P., Athanassiou C.G., Sarlis G.P., Petrović O., Niketić M. and Veroniki M.A. 2004. A Survey of Aphid Parasitoids (Hymenoptera: Braconidae: Aphidiinae) of Southeastern Europe and Their Aphid-Plant Associations. Applied Entomology and Zoology, 39 (3), 527-563.
- Kovar I. 2007. Coccinellidae, pp. 568-631. In I. Löbl and A. Smetana (ed.): Catalogue of Palearctic Coleoptera, Vol. 4. Stenstrup: Apollo Books, 935 pp.
- Kök Ş., Kasap İ. and Özdemir I. 2016. Aphid (Hemiptera: Aphididae) Species Determined in Çanakkale Province with a New Record for the Aphid Fauna of Turkey. Turkish Journal of Entomology, 40 (4), 397-412.
- Larson D.J. 2013. Key to Lady Beetles (Coleoptera: Coccinellidae) of Saskatchewan, [http://www.entsocsask.ca/documents/insect\\_lists/Coccinellidae\\_key.pdf](http://www.entsocsask.ca/documents/insect_lists/Coccinellidae_key.pdf) (Date accessed: June 2017).
- Magro A., Lecompte E., Magne F., Hemptinne J.L. and Crouau-Roy B. 2010. Phylogeny of Ladybirds (Coleoptera: Coccinellidae): Are the Subfamilies Monophyletic?. Molecular Phylogenetics and Evolution, 54, 833-848.

- Ölmez S. and Ulusoy M.R. 2003. A survey of Aphid Parasitoids (Hymenoptera: Braconidae: Aphidiinae) in Diyarbakır, Turkey. *Phytoparasitica*, 31 (5), 524-528.
- Poorani J. 2004. An Annotated Checklist of the Coccinellidae (Coleoptera) of Indian Subregion, (Web page: [http://www.angelfire.com/bug2/j\\_poorani/checklist.pdf](http://www.angelfire.com/bug2/j_poorani/checklist.pdf)) (Date accessed: June 2017).
- Posada-Flórez F.J., Téllez-Farfán L., Simbaqueba R. and Serna F. 2014. Survey and Bionomic Observations of *Siphaflava* Damaging Kikuyo Grass in Two Colombian Sites. *Revista U.D.C.A Actualidad and Divulgación Científica*, 17 (1), 95-104.
- Sharkey M.J. and Wharton R.A. 1997. Morphology and Terminology. In: R. A. Wharton, P. M. Marsh, and M. J. Sharkey (eds.), *Manual of the New World Genera of the Family Braconidae (Hymenoptera). Special Publication 1. International Society of Hymenopterists*, Washington, DC, 19-37 pp.
- Smith C.F. 1944. The Aphidiinae of North America. *Ohio State University Contributions in Zoology and Entomology Entstitute*, 6, 1-154.
- Starý P. 1970. Biology of Aphid Parasites (Hymenoptera: Aphidiidae) with respect to Integrated Control. *Series Entomologica*, Vol. 6. Dr. W. Junk, the Hague, the Netherlands, 656 pp.
- Starý P. 1988. Aphidiidae. In: Minks, A. K. and Harrewijn, P. (eds.), *Aphids: Their Biology, Enemies and Control*. vol. 2B. Elsevier, Amsterdam, the Netherlands. 171-184 pp.
- Starý P. 2006. Aphid parasitoids of the Czech Republic (Hymenoptera: Braconidae: Aphidiinae). *Academia, Praha*, 292 pp.
- Şahbaz A. and Uysal M. 2006. Konya İlinde Kavaklarda Beslenen Yaprakbitlerinin (Homoptera: Aphididae) Predatör ve Parazitoitleri. *Selçuk Üniversitesi Ziraat Fakültesi Dergisi*, 20 (38), 119-125.
- Tezcan S. and Uygun N. 2003. İzmir ve Manisa Yöresi Ekolojik Kiraz Üretim Bahçelerinde Saptanan Coccinellidae (Coleoptera) Türleri Üzerinde Bir Değerlendirme. *Türkiye Entomoloji Dergisi*, 27 (1), 73-79.
- Tomanović Ž. and Kavallieratos N.G. 2002. Two New Aphidiine Wasps (Hymenoptera: Braconidae: Aphidiinae) from the Southeastern Europe. *Reichenbachia*, 34, 341-345.
- Tomanović Ž. and Starý P. 2001. *Aphidius linosiphonis* sp.n. (Hymenoptera: Braconidae: Aphidiinae), a New Member of the Aphid Parasitoid Guild Associated with *Galium*. *Zootaxa*, 6, 1-4.
- Tomanović Ž., Beyarslan A., Çetin Erdoğan Ö. and Žikić V. 2008. New Records of Aphid Parasitoids (Hymenoptera, Braconidae, Aphidiinae) from Turkey. *Periodicum Biologorum*, 110 (4), 335-338.
- Tomanović Ž., Kavallieratos N.G., Athanassiou C.G. and Petrović O. 2003b. A New *Praon* Species (Hymenoptera: Braconidae: Aphidiinae) of the *Uroleucon* Parasitoid Complex from the Mediterranean Area. *Phytoparasitica*, 31, 19-26.
- Tomanović Ž., Kavallieratos N.G., Athanassiou C.G., Stanisljević L.Z. 2003a. A Review of the West Palaearctic Aphidiine (Hymenoptera: Braconidae: Aphidiinae) Parasitic

Aphidophagous coccinellid and parasitoid species determined in the Çanakkale Province with a new record for the parasitoid fauna of Turkey

on *Uroleucon* spp. with a Description of a New Species. Annales Societe Entomologique de France, 39, 343-353.

Tomanović Ž., Starý P. and Petrović O. 2002. *Monoctonus leclanti* sp.n. (Hymenoptera: Braconidae: Aphidiinae) From High Montane Areas of Southeastern Europe and key to Related Species. Entomologica Fennica, 13, 159-162.

Uygun N. 1981. Türkiye Coccinellidae (Col.) Faunası Üzerine Taksonomik Araştırmalar. Çukurova Üniversitesi Ziraat Fakültesi Yayınları, 157, 110 pp.

van Emden H.E. 1995. Host-Plant-Aphidophaga Interactions. Agriculture, Ecosystems and Environment, 52 (1), 3-11.

van Emden H.F., Harrington R. 2007. Aphids as Crop Pests. CABI Publishing, London, 717 pp.

Yousof D.E., Mahmoud M.E.E. and Mohamed A.H. 2013. Prospects of Biological Control of Daten Palm Green Pit Scale Insect *Asterolecanium phoenicis* Rao (Homoptera: Asterolecaniidae) in Sudan. Persian Gulf Crop Protection, 2 (2), 42-48.

Yu D.S., van Achterberg K. and Horstmann K. 2012. World Ichneumonoidea 2011, Taxapad 2012, (Web page: <http://www.taxapad.com/global.php>) (Date accessed: September 2016).

Bitki Koruma Bülteni Yayın Kurulu; makalelerin incelenmesi ve değerlendirilmesine katkıda bulunan aşağıda isimleri verilen konu uzmanlarına teşekkür eder.

**AKYOL, Yrd. Doç. Dr. Mustafa**-Manisa Celal Bayar Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Manisa

**ALTIN, Yrd. Doç. Dr. Nedim**-Düzce Üniversitesi, Ziraat ve Doğa Bilimleri Fakültesi, Bitki Koruma Bölümü, Düzce

**ALTINOK, Doç. Dr. Hacer Handan**-Erciyes Üniversitesi, Seyrani Ziraat Fakültesi, Bitki Koruma Bölümü, Kayseri.

**ANLAŞ, Doç. Dr. Sinan**-Manisa Celal Bayar Üniversitesi, Alaşehir Meslek Yüksekokulu, Alaşehir, Manisa

**ARAZ, Dr. Aydan**-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara

**ARSLAN, Prof. Dr. Ümit**-Uludağ Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Bursa

**ASAV, Dr. Ünal**-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara

**BABAROĞLU, Dr. Numan**-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara

**BAQUERO, Prof. Dr. Enrique**- University of Navarra, Department of Zoology and Ecology, Spain

**BİRGÜCÜ, Doç. Dr. Ali Kemal**-Süleyman Demirel Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Isparta

**BİRGÜCÜ, Doç. Dr. Ali Kemal**-Süleyman Demirel Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Isparta

**BOZKURT, Dr. Vildan**-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara

**CANIHOŞ, Dr. Ercan**-Biyolojik Mücadele Araştırma Enstitüsü Müdürlüğü, Adana

**ÇETİNTAŞ, Prof. Dr. Ramazan**-Sütçü İmam Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Kahramanmaraş

**ÇOBANOĞLU, Prof. Dr. Sultan**-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ankara

**DEMİRCİ, Prof. Dr. Fikret**-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ankara

**DESEN KÖYCÜ, Yrd. Doç. Dr. Nagehan**-Namık Kemal Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Tekirdağ

- DİLER, Zir. Yük. Müh. Heval-Zirai Mücadele Merkez Araştırma Enstitüsü**  
Müdürlüğü, Ankara
- DOĞAN, Prof. Dr. Salih-Erzincan Üniversitesi, Fen Fakültesi, Biyoloji Bölümü,**  
Erzincan
- ERDOĞAN ÇETİN, Prof. Dr. Özlem-Trakya Üniversitesi, Fen-Edebiyat**  
Fakültesi, Biyoloji Bölümü, Edirne
- ERDOĞAN, Doç. Dr. Oktay-Nevşehir Hacı Bektaş Veli Üniversitesi Mühendislik-**  
Mimarlık Fakültesi Biyosistem Mühendisliği Bölümü, Nevşehir
- ERKO, Dr. Fatma Dolunay-Tarımsal Araştırmalar ve Politikalar Genel**  
Müdürlüğü, Bitki Sağlığı Daire Başkanlığı, Ankara
- GÖZCÜ, Zir. Yük. Müh. Doğan-Doğu Akdeniz Geçit Kuşağı Tarımsal Araştırma**  
Enstitüsü Müdürlüğü, Kahramanmaraş
- GÜNAÇTI, Dr. Hale-Biyolojik Mücadele Araştırma Enstitüsü Müdürlüğü, Adana**
- GÜNER, Dr. Üftade-Zirai Mücadele Araştırma Enstitüsü Müdürlüğü, Bornova-**  
İzmir
- GÜVEN, Zir. Yük. Müh. Bilgin-Bornova Zirai Mücadele Araştırma Enstitüsü**  
Müdürlüğü, İzmir
- HEKİMİHAN, Dr. Hakan-Ege Tarımsal Araştırma Enstitüsü Müdürlüğü, İzmir**
- İŞİK, Prof. Dr. Doğan-Erciyes Üniversitesi, Ziraat Fakültesi, Bitki Koruma**  
Bölümü, Kayseri
- İNANÇ Denizhan, Yrd. Doç. Dr. Evsel-Van Yüzüncü Yıl Üniversitesi, Ziraat**  
Fakültesi, Bitki Koruma Bölümü, Van
- KAPLAN, Yrd. Doç. Dr. Cevdet-Siirt Üniversitesi, Ziraat Fakültesi, Bitki Koruma**  
Bölümü, Siirt
- KARACA, Prof. Dr. Gürsel-Süleyman Demirel Üniversitesi, Ziraat Fakültesi,**  
Bitki Koruma Bölümü, Isparta
- KARACAOĞLU, Dr. Mehmet-Biyolojik Mücadele Araştırma Enstitüsü**  
Müdürlüğü, Adana
- KARAKAYA, Prof. Dr. Aziz-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma**  
Bölümü, Ankara
- KARAKOÇ, Yrd. Doç. Dr. Ömer Cem-Çankırı Karatekin Üniversitesi, Yapraklı**  
Meslek Yüksekokulu, Bitkisel ve Hayvansal Üretim Bölümü, Çankırı
- KAYA, Zir. Yük. Müh. Yalçın-Tokat Orta Karadeniz Geçit Kuşağı Tarımsal**  
Araştırma Enstitüsü Müdürlüğü, Tokat

- KEÇECİ, Yrd. Doç. Dr. Mehmet-İnönü** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Malatya
- KILINÇ, Yrd. Doç. Dr. Özgür Kıvılcım-Niğde** Ömer Halisdemir Üniversitesi, Bitkisel Üretim Ve Teknolojileri Bölümü, Niğde
- KURT, Prof. Dr. Şener-Mustafa Kemal** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Hatay
- KUTLUK YILMAZ, Doç. Dr. Nazlı Dide-Ondokuz** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Samsun
- MERMER DOĞU, Zir. Yük. Müh. Duygu-Zirai Mücadele Merkez Araştırma** Enstitüsü Müdürlüğü, Ankara
- MUTLU, Yrd. Doç. Dr. Çetin-Abant İzzet Baysal** Üniversitesi Ziraat ve Doğa Bilimleri Fakültesi, Bolu
- OKSAL, Yrd. Doç. Dr. Hatice Diğdem-İnönü** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Malatya
- OLHAN, Prof. Dr. Emine-Ankara** Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Ankara
- ÖZÇELİK, Prof. Dr. Ahmet-Ankara** Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Ankara
- ÖZDEN DEMİR, Doç. Dr. Emine-Düzce** Üniversitesi, Ziraat ve Doğa Bilimleri Fakültesi, Tarla Bitkileri Bölümü, Düzce
- ÖZER, Prof. Dr. Nuray-Namık Kemal** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Tekirdağ
- ÖZPINAR, Prof. Dr. Ali-Çanakkale Onsekiz Mart** Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Çanakkale
- ÖZSEMERÇİ, Dr. Fatma-Bornova Zirai Mücadele Araştırma Enstitüsü** Müdürlüğü, İzmir
- PER, Yrd. Doç. Dr. Sedat-Bozok** Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Yozgat
- SERİM, Dr. Tansel-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü,** Ankara
- SEVGİLİ, Prof. Dr. Hasan-Ordu** Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Ordu
- SHAYANMEHR, Dr. Masoumeh-Sari University of Agricultural Sciences and** Natural Resources, Iran
- ŞABANOĞLU, Dr. Burcu-Hacettepe** Üniversitesi, Fen Fakültesi, Biyoloji Bölümü, Ankara

- ŞENYÜZ, Yrd. Doç. Dr. Yakup**-Dumlupınar Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Kütahya
- TARLA, Doç. Dr. Şener**-Uşak Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Uşak.
- TUNCA, Doç. Dr. Hilal**-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ankara
- TUNCA, Yrd. Doç. Dr. Hilal**-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ankara
- TÜLEK, Zir. Yük. Müh. Senem**-Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara
- TÜRKKAN, Doç. Dr. Muharrem**-Ordu Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ordu
- TÜRKSEVEN, Dr. Süleyman**-Ege Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, İzmir
- UĞUR, Prof. Dr. Avni**-Ankara Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Ankara
- URHAN, Prof. Dr. Raşit**-Pamukkale Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Denizli
- UYSAL, Dr. Osman**-Alata Bahçe Kültürleri Araştırma Enstitüsü Müdürlüğü, Mersin
- YANAR, Doç. Dr. Dürdane**-Gaziosmanpaşa Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Tokat
- YANAR, Prof. Dr. Yusuf**-Gaziosmanpaşa Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Tokat
- YILAR, Yrd. Doç. Dr. Melih**-Gaziosmanpaşa Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Tokat
- YOLDAŞ, Prof. Dr. Zeynep**-Ege Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, İzmir
- YÜCEL, Doç. Dr. Seral**-Biyolojik Mücadele Araştırma Enstitüsü Müdürlüğü, Adana
- ZEYBEKOĞLU, Prof. Dr. Ünal**-Ondokuz Mayıs Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, Samsun

The Editorial Board thanks scientific advisory board listed below for their contributions to redaction.

**AKYOL, Yrd. Doç. Dr. Mustafa-**Manisa Celal Bayar University, Faculty of Arts and Science, Department of Biological Sciences, Manisa

**ALTIN, Yrd. Doç. Dr. Nedim-**Düzce University, Faculty of Agriculture and Natural Sciences, Department of Plant Protection, Düzce

**ALTINOK, Doç. Dr. Hacer Handan-**Erciyes University, Seyrani Agricultural Faculty, Department of Plant Protection, Kayseri

**ANLAŞ, Doç. Dr. Sinan-**Manisa Celal Bayar University, Alaşehir Vocational School, Alaşehir, Manisa

**ARAZ, Dr. Aydan-**Directorate of Plant Protection Central Research Institute, Ankara

**ARSLAN, Prof. Dr. Ümit-**Uludağ University, Faculty of Agriculture, Department of Plant Protection, Bursa

**ASAV, Dr. Ünal-**Directorate of Plant Protection Central Research Institute, Ankara

**BABAROĞLU, Dr. Numan-**Directorate of Plant Protection Central Research Institute, Ankara

**BAQUERO, Prof. Dr. Enrique-**University of Navarra, Department of Zoology and Ecology, Spain

**BİRGÜCÜ, Doç. Dr. Ali Kemal-**Süleyman Demirel University, Faculty of Agriculture, Department of Plant Protection, Isparta

**BİRGÜCÜ, Doç. Dr. Ali Kemal-**Süleyman Demirel University, Faculty of Agriculture, Department of Plant Protection, Isparta

**BOZKURT, Dr. Vildan-**Directorate of Plant Protection Central Research Institute, Ankara

**CANIHOŞ, Dr. Ercan-**Directorate of Biological Control Research Institute, Adana

**ÇETİNTAŞ, Prof. Dr. Ramazan-**Sütçü İmam University, Faculty of Agriculture, Department of Plant Protection, Kahramanmaraş

**ÇOBANOĞLU, Prof. Dr. Sultan-**Ankara University, Faculty of Agriculture, Department of Plant Protection, Ankara

**DEMİRCİ, Prof. Dr. Fikret-**Ankara University, Faculty of Agriculture, Department of Plant Protection, Ankara

**DESEN KÖYCÜ, Yrd. Doç. Dr. Nagehan-**Namık Kemal University, Faculty of Agriculture, Department of Plant Protection, Tekirdağ

**DİLER, Zir. Yük. Müh. Heval**-Directorate of Plant Protection Central Research Institute, Ankara

**DOĞAN, Prof. Dr. Salih**-Erzincan University, Faculty of Science, Department of Biological Sciences, Erzincan

**ERDOĞAN ÇETİN, Prof. Dr. Özlem**-Trakya University, Faculty of Arts and Science, Department of Biological Sciences, Edirne

**ERDOĞAN, Doç. Dr. Oktay**-The University of Nevşehir Hacı Bektaş Veli Engineering - Architecture Faculty Department of Biosystem Engineering, Nevşehir.

**ERKOL, Dr. Fatma Dolunay**-General Directorate of Agricultural Research and Policies, Ankara

**GÖZCÜ, Zir. Yük. Müh. Doğan**- East Mediterranean Transitional Zone Agricultural Research of Institute

**GÜNAÇTI, Dr. Hale**- Directorate of Biological Control Research Institute, Adana

**GÜNER, Dr. Üftade**- Directorate of Plant Protection Research Institute Bornova, İzmir

**GÜVEN, Zir. Yük. Müh. Bilgin**- Directorate of Plant Protection Research Institute Bornova, İzmir

**HEKİMHAN, Dr. Hakan**- Aegean Agricultural Research Institute, İzmir

**IŞIK, Prof. Dr. Doğan**-Erciyes University, Faculty of Agriculture, Department of Plant Protection, Kayseri

**İNANÇ Denizhan, Yrd. Doç. Dr. Evsel**-Van Yüzüncü Yıl University, Faculty of Agriculture, Department of Plant Protection, Van

**KAPLAN, Yrd. Doç. Dr. Cevdet**-Siirt University, Faculty of Agriculture, Department of Plant Protection, Siirt

**KARACA, Prof. Dr. Gürsel**- Süleyman Demirel University, Faculty of Agriculture, Department of Plant Protection, Isparta

**KARACAOĞLU, Dr. Mehmet**- Directorate of Biological Control Research Institute, Adana

**KARAKAYA, Prof. Dr. Aziz**- Ankara University, Faculty of Agriculture, Department of Plant Protection, Ankara

**KARAKOÇ, Yrd. Doç. Dr. Ömer Cem**- Cankiri Karatekin University, Yapraklı Vocational School, Department of Crop and Animal Protection Çankırı

**KAYA, Zir. Yük. Müh. Yalçın**- Middle Black Sea Transitional Zone Agricultural Research Institute, Tokat

- KEÇECİ, Yrd. Doç. Dr. Mehmet-** İnönü University, Faculty of Agriculture, Department of Plant Protection, Malatya
- KILINÇ, Yrd. Doç. Dr. Özgür Kıvılcım-**Niğde Ömer Halisdemir University, Faculty of Agricultural Sciences and Technologies, Plant Production and Technologies, Niğde
- KURT, Prof. Dr. Şener-**Mustafa Kemal University, Faculty of Agriculture, Department of Plant Protection, Hatay
- KUTLUK YILMAZ, Doç. Dr. Nazlı Dide-**Ondokuz Mayıs University, Faculty of Agriculture, Department of Plant Protection, Samsun
- MERMER DOĞU, Zir. Yük. Müh. Duygu-**Directorate of Plant Protection Central Research Institute, Ankara
- MUTLU, Yrd. Doç. Dr. Çetin-**Abant İzzet Baysal University, Faculty of Agriculture and Natural Sciences, Department of Plant Protection, Bolu
- OKSAL, Yrd. Doç. Dr. Hatice Diğdem-**İnönü University, Faculty of Agriculture, Department of Plant Protection, Malatya
- OLHAN, Prof. Dr. Emine-**Ankara University, Faculty of Agriculture, Department of Agricultural Economics, Ankara
- ÖZÇELİK, Prof. Dr. Ahmet-**Ankara University, Faculty of Agriculture, Department of Agricultural Economics, Ankara Ankara
- ÖZDEN DEMİR, Doç. Dr. Emine-**Düzce University, Faculty of Agriculture and Natural Sciences, Department of Field Crop, Düzce
- ÖZER, Prof. Dr. Nuray-** Namık Kemal University, Faculty of Agriculture, Department of Plant Protection, Tekirdağ
- ÖZPINAR, Prof. Dr. Ali-** Çanakkale Onsekiz Mart University, Faculty of Agriculture, Department of Plant Protection, Çanakkale
- ÖZSEMERÇİ, Dr. Fatma-** Directorate of Plant Protection Research Institute Bornova, İzmir
- PER, Yrd. Doç. Dr. Sedat-**Bozok University, Faculty of Arts and Science, Department of Biological Sciences, Yozgat
- SERİM, Dr. Tansel-** Directorate of Plant Protection Central Research Institute, Ankara
- SEVGİLİ, Prof. Dr. Hasan-**Ordu University, Faculty of Arts and Science, Department of Biological Sciences, Ordu
- SHAYANMEHR, Dr. Masoumeh-** Sari University of Agricultural Sciences and Natural Resources, Iran

**ŞABANOĞLU, Dr. Burcu**-Hacettepe University, Faculty of Science, Department of Biological Sciences, Ankara

**ŞENYÜZ, Yrd. Doç. Dr. Yakup**-Dumlupınar University, Faculty of Arts and Science, Department of Biological Sciences, Kütahya

**TARLA, Doç. Dr. Şener**-Uşak University, Faculty of Agriculture, Department of Plant Protection, Uşak.

**TUNCA, Doç. Dr. Hilal**- Ankara University, Faculty of Agriculture, Department of Plant Protection, Ankara

**TÜLEK, Zir. Yük. Müh. Senem**- Directorate of Plant Protection Central Research Institute, Ankara

**TÜRKKAN, Doç. Dr. Muharrem**-Ordu University, Faculty of Agriculture, Department of Plant Protection, Ordu

**TÜRKSEVEN, Dr. Süleyman**- Ege University, Faculty of Agriculture, Department of Plant Protection, İzmir

**UĞUR, Prof. Dr. Avni**- Ankara University, Faculty of Agriculture, Department of Plant Protection, Ankara

**URHAN, Prof. Dr. Raşit**-Pamukkale University, Faculty of Arts and Science, Department of Biological Sciences, Denizli

**UYSAL, Dr. Osman**-Alata Horticultural Research Institute, Mersin

**YANAR, Doç. Dr. Dürdane**- Gaziosmanpaşa University, Faculty of Agriculture, Department of Plant Protection, Tokat

**YANAR, Prof. Dr. Yusuf**- Gaziosmanpaşa University, Faculty of Agriculture, Department of Plant Protection, Tokat

**YILAR, Yrd. Doç. Dr. Melih**- Gaziosmanpaşa University, Faculty of Agriculture, Department of Plant Protection, Tokat

**YOLDAŞ, Prof. Dr. Zeynep**- Ege University, Faculty of Agriculture, Department of Plant Protection, İzmir

**YÜCEL, Doç. Dr. Seral**- Directorate of Biological Control Research Institute, Adana

**ZEYBEKOĞLU, Prof. Dr. Ünal**- Ondokuz Mayıs University, Faculty of Agriculture, Department of Plant Protection, Samsun